

A KOHNO–DRINFELD THEOREM FOR QUANTUM WEYL GROUPS

VALERIO TOLEDANO LAREDO

ABSTRACT. Let $\mathfrak{g}$ be a complex, simple Lie algebra with Cartan subalgebra $\mathfrak{h}$ and Weyl group W . In [MTL], we introduced a new, W –equivariant flat connection on $\mathfrak{h}$ with simple poles along the root hyperplanes and values in any finite–dimensional $\mathfrak{g}$ –module V . It was conjectured in [TL] that its monodromy is equivalent to the quantum Weyl group action of the generalised braid group of type $\mathfrak{g}$ on V obtained by regarding the latter as a module over the quantum group $U_{\hbar}\mathfrak{g}$. In this paper, we prove this conjecture for $\mathfrak{g} = \mathfrak{sl}_n$.

1. INTRODUCTION

One of the many virtues of quantum groups is their ability to describe the monodromy of certain first order systems of Fuchsian PDEs. If $U_{\hbar}\mathfrak{g}$ is the Drinfeld–Jimbo quantum group of the complex, simple Lie algebra $\mathfrak{g}$, the universal R –matrix of $U_{\hbar}\mathfrak{g}$ yields a representation of Artin’s braid group on n strings B_n on the n –fold tensor product $V^{\otimes n}$ of any finite–dimensional $U_{\hbar}\mathfrak{g}$ –module V . A fundamental, and paradigmatic result of Kohno and Drinfeld establishes the equivalence of this representation with the monodromy of the Knizhnik–Zamolodchikov equations for $\mathfrak{g}$ with values in $V^{\otimes n}$ [Dr3, Dr4, Dr5, Ko1]. Lusztig, and independently Kirillov–Reshetikhin and Soibelman realised that $U_{\hbar}\mathfrak{g}$ also yields representations of another braid group, namely the generalised braid group $B_{\mathfrak{g}}$ of Lie type $\mathfrak{g}$ [Lu1, KR, So]. Whereas the R –matrix representation is a deformation of the natural action of the symmetric group $\mathfrak{S}_n$ on n –fold tensor products, these representations of $B_{\mathfrak{g}}$ deform the action of (a finite extension of) the Weyl group W of $\mathfrak{g}$ on any finite–dimensional $\mathfrak{g}$ –module V .

The aim of this paper is to show that these quantum Weyl group representations describe the monodromy of the flat connection introduced in [MTL] and, independently, in [FMTV]. More precisely, realise $B_{\mathfrak{g}}$ as the fundamental group of the orbit space $\mathfrak{h}_{\text{reg}}/W$ of the set of regular elements of a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ under the action of W [Br]. Then, one can define a flat vector bundle $(\mathcal{V}, \nabla_{\kappa})$ with fibre V over $\mathfrak{h}_{\text{reg}}/W$ [MTL]. The connection ∇_{κ} depends upon a parameter $\hbar \in \mathbb{C}$ and it was conjectured in [TL] that, when $\hbar$ is regarded as a formal variable, its monodromy is equivalent to the quantum Weyl group action of $B_{\mathfrak{g}}$ on V . This conjecture was checked in [TL] for a number of pairs $(\mathfrak{g}, V)$ including vector representations of classical Lie algebras and adjoint representations of simple Lie algebras.

In the present paper, we prove this conjecture for $\mathfrak{g} = \mathfrak{sl}_n$, so that $B_{\mathfrak{g}} = B_n$. The proof relies on the Kohno–Drinfeld theorem for $U_{\hbar}\mathfrak{sl}_k$ via the use of the dual pair $(\mathfrak{gl}_k, \mathfrak{gl}_n)$. Our main observation is that the duality between $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ derived from their joint action on the space $\mathcal{M}_{k,n}$ of $k \times n$ matrices exchanges ∇_{κ} for $\mathfrak{sl}_n$ and the Knizhnik–Zamolodchikov connection for $\mathfrak{sl}_k$, thus acting as a simple–minded integral transform. This shows the equivalence of the monodromy representation of ∇_{κ} for $\mathfrak{sl}_n$ with a suitable R –matrix representation for $U_{\hbar}\mathfrak{sl}_k$. The proof is completed by noting that the duality between $U_{\hbar}\mathfrak{gl}_k$ and $U_{\hbar}\mathfrak{gl}_n$ exchanges the R –matrix representation of $U_{\hbar}\mathfrak{sl}_k$ with the quantum Weyl group representation of $U_{\hbar}\mathfrak{sl}_n$.

This may be schematically summarised by the following diagram

$$\begin{array}{ccc}
 \nabla_{\text{KZ}}, \mathfrak{sl}_k & \xleftrightarrow{\mathcal{M}_{k,n}} & \nabla_{\kappa}, \mathfrak{sl}_n \\
 \uparrow KD & & \uparrow \\
 R^{\vee}, U_{\hbar} \mathfrak{sl}_k & \xleftrightarrow{\mathcal{M}_{k,n}^{\hbar}} & W_{\hbar}, U_{\hbar} \mathfrak{sl}_n
 \end{array}$$

The structure of the paper is as follows. In section 2, we give the construction of the connection ∇_{κ} following [MTL]. We show in section 3 that the duality between $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ identifies the Knizhnik–Zamolodchikov connection for n -fold tensor products of symmetric powers of the vector representation of $\mathfrak{sl}_k$ and the connection ∇_{κ} for $\mathfrak{sl}_n$. In section 4 we recall the definition of the Drinfeld–Jimbo quantum groups $U_{\hbar} \mathfrak{gl}_k$ and $U_{\hbar} \mathfrak{gl}_n$ and, in section 5 show how they jointly act on the quantum $k \times n$ matrix space $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$. The corresponding R -matrix and quantum Weyl group representations of B_n on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ are shown to coincide in section 6. Section 7 contains our main result.

Acknowledgements. This paper was begun at the Research Institute for Mathematical Sciences of Kyoto University. I am very grateful to M. Kashiwara for his invitation to spend the summer of 1999 at RIMS and to RIMS for its hospitality and financial support. During my stay, I greatly benefitted from very stimulating and informative discussions with M. Kashiwara and B. Feigin. I also wish to express my gratitude to A. D’Agnolo, P. Baumann, B. Enriquez, J. Millson, R. Rouquier and P. Schapira for innumerable, useful and friendly conversations.

2. FLAT CONNECTIONS ON $\mathfrak{h}_{\text{reg}}$

The results in this section are due to J. Millson and the author [MTL]. They were obtained independently by De Concini around 1995 (unpublished). Let $\mathfrak{g}$ be a complex, simple Lie algebra with Cartan subalgebra $\mathfrak{h}$ and root system $R \subset \mathfrak{h}^*$. Let $\mathfrak{h}_{\text{reg}} = \mathfrak{h} \setminus \bigcup_{\alpha \in R} \text{Ker}(\alpha)$ be the set of regular elements in $\mathfrak{h}$ and V a finite-dimensional $\mathfrak{g}$ -module. We shall presently define a flat connection on the topologically trivial vector bundle $\mathfrak{h}_{\text{reg}} \times V$ over $\mathfrak{h}_{\text{reg}}$. We need for this purpose the following simple flatness criterion due to Kohno [Ko2]. Let B be a complex, finite-dimensional vector space and $\mathcal{A} = \{H_i\}_{i \in I}$ a finite collection of hyperplanes in B determined by the linear forms $\phi_i \in B^*$, $i \in I$.

Lemma 2.1. *Let F be a finite-dimensional vector space and $\{r_i\} \subset \text{End}(F)$ a family indexed by I . Then,*

$$\nabla = d - \sum_{i \in I} \frac{d\phi_i}{\phi_i} r_i \quad (2.1)$$

defines a flat connection on $(B \setminus \mathcal{A}) \times F$ iff, for any subset $J \subseteq I$ maximal for the property that $\bigcap_{j \in J} H_j$ is of codimension 2, the following relations hold for any $j \in J$

$$[r_j, \sum_{j' \in J} r_{j'}] = 0 \quad (2.2)$$

For any $\alpha \in R$, choose root vectors $e_\alpha \in \mathfrak{g}_\alpha, f_\alpha \in \mathfrak{g}_{-\alpha}$ such that $[e_\alpha, f_\alpha] = h_\alpha = \alpha^\vee$ and let

$$\kappa_\alpha = \frac{\langle \alpha, \alpha \rangle}{2} (e_\alpha f_\alpha + f_\alpha e_\alpha) \in U\mathfrak{g} \quad (2.3)$$

be the truncated Casimir operator of the $\mathfrak{sl}_2(\mathbb{C})$ -subalgebra of $\mathfrak{g}$ spanned by $e_\alpha, h_\alpha, f_\alpha$. Note that κ_α does not depend upon the particular choice of e_α and f_α and that $\kappa_{-\alpha} = \kappa_\alpha$.

Theorem 2.2. *The one-form*

$$\nabla_\kappa^h = d - h \sum_{\alpha > 0} \frac{d\alpha}{\alpha} \kappa_\alpha = d - \frac{h}{2} \sum_{\alpha \in R} \frac{d\alpha}{\alpha} \kappa_\alpha \quad (2.4)$$

defines, for any $h \in \mathbb{C}$, a flat connection on $\mathfrak{h}_{\text{reg}} \times V$.

PROOF. By lemma 2.1, we must prove that for any rank 2 subsystem $R_0 \subseteq R$, the following holds for any $\alpha \in R_0^+ = R_0 \cap R^+$

$$[\kappa_\alpha, \sum_{\beta \in R_0^+} \kappa_\beta] = 0 \quad (2.5)$$

This may be proved by an explicit computation by considering in turn the cases where R_0 is of type $A_1 \times A_1, A_2, B_2$ or G_2 but is more easily settled by the following elegant observation of A. Knutson [Kn]. Let $\mathfrak{g}_0 \subseteq \mathfrak{g}$ be the semi-simple Lie algebra with root system R_0 , $\mathfrak{h}_0 \subset \mathfrak{h}$ its Cartan subalgebra and $C_0 \in Z(U\mathfrak{g}_0)$ its Casimir operator. Then, $\sum_{\beta \in R_0^+} \kappa_\beta - C_0$ lies in $U\mathfrak{h}_0$ so that (2.5) holds since κ_α commutes with $\mathfrak{h}_0$ ■

Let G be the complex, connected and simply-connected Lie group with Lie algebra $\mathfrak{g}$, T its torus with Lie algebra $\mathfrak{h}$, $N(T) \subset G$ the normaliser of T and $W = N(T)/T$ the Weyl group of G . Let $B_{\mathfrak{g}} = \pi_1(\mathfrak{h}_{\text{reg}}/W)$ be the generalised braid group of type $\mathfrak{g}$ and $\sigma : B_{\mathfrak{g}} \rightarrow N(T)$ a homomorphism compatible with

$$\begin{array}{ccc} B_{\mathfrak{g}} & \xrightarrow{\sigma} & N(T) \\ & \searrow & \downarrow \\ & & W \end{array} \quad (2.6)$$

[Ti]. We regard $B_{\mathfrak{g}}$ as acting on V via σ . Let $\widetilde{\mathfrak{h}_{\text{reg}}} \xrightarrow{p} \mathfrak{h}_{\text{reg}}$ be the universal cover of $\mathfrak{h}_{\text{reg}}$ and $\mathfrak{h}_{\text{reg}}/W$.

Proposition 2.3. *The one-form $p^* \nabla_\kappa^h$ defines a $B_{\mathfrak{g}}$ -equivariant flat connection on $\widetilde{\mathfrak{h}_{\text{reg}}} \times V = p^*(\mathfrak{h}_{\text{reg}} \times V)$. It therefore descends to a flat connection on the vector bundle*

$$\begin{array}{ccc} V & \longrightarrow & \widetilde{\mathfrak{h}_{\text{reg}}} \times_{B_{\mathfrak{g}}} V \\ & & \downarrow \\ & & \mathfrak{h}_{\text{reg}}/W \end{array} \quad (2.7)$$

which is reducible with respect to the weight space decomposition of V and unitary if $h \in i\mathbb{R}$.

PROOF. The action of $B_{\mathfrak{g}}$ on $\Omega^\bullet(\widetilde{\mathfrak{h}_{\text{reg}}}, V) = \Omega^\bullet(\widetilde{\mathfrak{h}_{\text{reg}}}) \otimes V$ is given by $\gamma \rightarrow (\gamma^{-1})^* \otimes \sigma(\gamma)$. Thus, if $\gamma \in B_{\mathfrak{g}}$ projects onto $w \in W$, we get using $p \cdot \gamma^{-1} = w^{-1} \cdot p$,

$$\gamma p^* \nabla_\kappa^h \gamma^{-1} = d - \frac{h}{2} \sum_{\alpha \in R} dp^* w \alpha / p^* w \alpha \otimes \sigma(\gamma) \kappa_\alpha \sigma(\gamma)^{-1} \quad (2.8)$$

Since $\kappa_\alpha = \frac{\langle \alpha, \alpha \rangle}{2}(e_\alpha f_\alpha + f_\alpha e_\alpha)$ is independent of the choice of the root vectors e_α, f_α , $\text{Ad}(\sigma(\gamma))\kappa_\alpha = \kappa_{w\alpha}$ and (2.8) is equal to $p^*\nabla_\kappa^h$ as claimed. $p^*\nabla_\kappa^h$ is flat by theorem 2.2, commutes with the fibrewise action of $\mathfrak{h}$ because each κ_α is of weight 0 and is unitary because the κ_α are self-adjoint ■

Thus, for any homomorphism $\sigma : B_{\mathfrak{g}} \rightarrow N(T)$ compatible with (2.6), proposition 2.3 yields a monodromy representation $\rho_h^\sigma : B_{\mathfrak{g}} \rightarrow GL(V)$ which permutes the weight spaces compatibly with W . By standard ODE theory, ρ_h^σ depends analytically on the complex parameter h and, when $h = 0$, is equal to the action of $B_{\mathfrak{g}}$ on V given by σ . We record for later use the following elementary

Proposition 2.4. *Let $\gamma \in B_{\mathfrak{g}} = \pi_1(\mathfrak{h}_{\text{reg}}/W)$ and $\tilde{\gamma} : [0, 1] \rightarrow \mathfrak{h}_{\text{reg}}$ be a lift of γ . Then,*

$$\rho_h^\sigma(\gamma) = \sigma(\gamma)\mathcal{P}(\tilde{\gamma}) \quad (2.9)$$

where $\mathcal{P}(\tilde{\gamma}) \in GL(V)$ is the parallel transport along $\tilde{\gamma}$ for the connection ∇_κ^h on $\mathfrak{h}_{\text{reg}} \times V$.

PROOF. Let $\tilde{\tilde{\gamma}} : [0, 1] \rightarrow \widetilde{\mathfrak{h}_{\text{reg}}}$ be a lift of γ and $\tilde{\gamma}$ so that $\tilde{\tilde{\gamma}}(1) = \gamma^{-1}\tilde{\tilde{\gamma}}(0)$. Then, since the connection on $p^*(\mathfrak{h}_{\text{reg}} \times V)$ is the pull-back of ∇_κ^h , and that on $(p^*(\mathfrak{h}_{\text{reg}} \times V))/B_{\mathfrak{g}}$ the quotient of $p^*\nabla_\kappa^h$, we find

$$\rho_h^\sigma(\gamma) = \mathcal{P}(\gamma) = \sigma(\gamma)\mathcal{P}(\tilde{\tilde{\gamma}}) = \sigma(\gamma)\mathcal{P}(\tilde{\gamma}) \quad (2.10)$$

■

By [Br], $B_{\mathfrak{g}}$ is presented on generators T_i , $i = 1 \dots n$ labelled by a choice of simple roots α_i of R with relations

$$T_i T_j T_i \dots = T_j T_i T_j \dots \quad (2.11)$$

for any $1 \leq i < j \leq n$ where each side of (2.11) has a number of factors equal to the order of $s_i s_j$ in W and $s_k \in W$ is the orthogonal reflection across the hyperplane $\text{Ker}(\alpha_k)$. T_i projects onto $s_i \in W$. An explicit choice of representatives of $T_1, \dots, T_n$ in $\pi_1(\mathfrak{h}_{\text{reg}}/W)$ may be given as follows. Let $t \in \mathfrak{h}_{\text{reg}}$ lie in the fundamental Weyl chamber so that $\langle t, \alpha \rangle > 0$ for any $\alpha \in R_+$. Note that for any simple root α_i , the intersection $t_{\alpha_i} = t - \frac{1}{2}\langle t, \alpha_i \rangle \alpha_i^\vee$ of the affine line $t + \mathbb{C} \cdot \alpha_i^\vee$ with $\text{Ker}(\alpha_i)$ does not lie in any other root hyperplane $\text{Ker}(\beta)$, $\beta \in R \setminus \{\alpha_i\}$. Indeed, if $\langle t_{\alpha_i}, \beta \rangle = 0$ then

$$2\langle t, \beta \rangle = \langle t, \alpha_i \rangle \langle \alpha_i^\vee, \beta \rangle = \langle t, \beta - s_i \beta \rangle \quad (2.12)$$

whence $\langle t, \beta \rangle = -\langle t, s_i \beta \rangle$, a contradiction since s_i permutes positive roots different from α_i . Let now D be an open disc in $t + \mathbb{C} \cdot \alpha_i^\vee$ of center t_{α_i} such that its closure $\overline{D}$ does not intersect any root hyperplane other than $\text{Ker}(\alpha_i)$. Consider the path $\gamma_i : [0, 1] \rightarrow t + \mathbb{C} \cdot \alpha_i^\vee$ from t to $s_i t$ determined by $\gamma_i|_{[0, 1/3] \cup [2/3, 1]}$ is affine and lies in $t + \mathbb{R} \cdot \alpha^\vee \setminus D$, $\gamma_i(1/3), \gamma_i(2/3) \in \partial \overline{D}$ and $\gamma_i|_{[1/3, 2/3]}$ is a semicircular arc in $\partial \overline{D}$, positively oriented with respect to the natural orientation of $t + \mathbb{C} \cdot \alpha_i$. Then, the image of γ_i in $\mathfrak{h}_{\text{reg}}/W$ is a representative of T_i in $\pi_1(\mathfrak{h}_{\text{reg}}/W, Wt)$ [Br].

3. KNIZHNIK–ZAMOLODCHIKOV EQUATIONS AND DUAL PAIRS

We show in this section that the joint action of $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ on the space $\mathcal{M}_{k,n}$ of $k \times n$ matrices identifies the connection ∇_κ for $\mathfrak{g} = \mathfrak{sl}_n$ and the Knizhnik–Zamolodchikov connection for $\mathfrak{sl}_k$.

Let $\mathcal{S}(\mathcal{M}_{k,n}^*) = \mathbb{C}[x_{11}, \dots, x_{kn}]$ be the algebra of polynomial functions on $\mathcal{M}_{k,n}$. The group $GL_k \times GL_n$ acts on $\mathcal{S}(\mathcal{M}_{k,n}^*)$ by

$$(g_k, g_n) p(x) = p(g_k^t x g_n) \quad (3.1)$$

and leaves the homogeneous components $\mathcal{S}^d(\mathcal{M}_{k,n}^*)$ of $\mathcal{S}(\mathcal{M}_{k,n}^*)$, $d \in \mathbb{N}$, invariant. The decomposition of $\mathcal{S}(\mathcal{M}_{k,n}^*)$ under $GL_k \times GL_n$ is well-known (see *e.g.*, [Zh, §132] which we follow closely or [Mc, §1.4]¹). Let N_k, N_n be the groups of $k \times k$ and $n \times n$ upper triangular unipotent matrices respectively.

Lemma 3.1.

$$\mathcal{S}(\mathcal{M}_{k,n}^*)^{N_k \times N_n} = \mathbb{C}[\Delta_1, \dots, \Delta_{\min(k,n)}] \quad (3.2)$$

where $\Delta_l(x) = \det(x_{ij})_{1 \leq i,j \leq l}$ is the l th principal minor of the matrix x .

PROOF. Assume for simplicity that $k \leq n$. Let $\mathcal{D} \subset \mathcal{S}(\mathcal{M}_{k,n}^*)$ be the subset of matrices x such that $\Delta_i(x) \neq 0$ for $i = 1 \dots k$. By the Gauss decomposition, any $x \in \mathcal{D}$ is conjugate under $N_k^t \times N_n$ to a unique $k \times n$ matrix $d(x)$ with the same principal minors as x , diagonal principal $k \times k$ block and the remaining columns equal to zero. Consider now the $k \times n$ matrix

$$m(x) = \begin{pmatrix} \Delta_1(x) & \Delta_2(x) & \Delta_3(x) & \cdots & \Delta_{k-1}(x) & \Delta_k(x) & 0 & \cdots & 0 \\ -1 & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ 0 & -1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & -1 & 0 & 0 & \cdots & 0 \end{pmatrix} \quad (3.3)$$

Since $\Delta_i(m(x)) = \Delta_i(x)$, $1 \leq i \leq k$, $m(x)$ is also conjugate to $d(x)$, and therefore to x , under $N_k^t \times N_n$. Thus, by density of $\mathcal{D}$, a polynomial $p \in \mathcal{S}(\mathcal{M}_{k,n}^*)$ is invariant under $N_k \times N_n$ iff it is a function of $m(x)$ and therefore iff it is a polynomial in $\Delta_1, \dots, \Delta_k$ ■

Let $\mathbb{Y}_p \subset \mathbb{N}^p$ be the set of Young diagrams with at most p rows. For $\lambda \in \mathbb{Y}_p$, set $|\lambda| = \sum_{i=1}^p \lambda_i$ and let $V_\lambda^{(p)}$ be the irreducible representation of $GL_p(\mathbb{C})$ of highest weight λ .

Theorem 3.2. As $GL_k \times GL_n$ -modules,

$$\mathcal{S}^d(\mathcal{M}_{k,n}^*) \cong \bigoplus_{\substack{\lambda \in \mathbb{Y}_{\min(k,n)}, \\ |\lambda|=d}} V_\lambda^{(k)} \otimes V_\lambda^{(n)} \quad (3.4)$$

PROOF. Assume again $k \leq n$ for simplicity. By lemma 3.1, the highest weight vectors for the action of $GL_k(\mathbb{C}) \times GL_n(\mathbb{C})$ on $\mathcal{S}(\mathcal{M}_{k,n}^*)$ are the polynomials in $\Delta_1, \dots, \Delta_k$ which are eigenvectors for the torus of $GL_k \times GL_n$. Since Δ_l is of weight $\varpi_l^{(k)} \oplus \varpi_l^{(n)}$, where $\varpi_l^{(p)}$ is the l th fundamental weight of GL_p , the highest weight vectors are the monomials $\Delta_1^{m_1} \cdots \Delta_k^{m_k}$ with corresponding pair of Young diagrams (λ, λ) where

$$\lambda = (m_1 + \cdots + m_k, m_2 + \cdots + m_k, \dots, m_k) \quad (3.5)$$

Thus, (3.4) holds since Δ_l is a homogeneous function of degree l ■

As a $\mathfrak{gl}_k$ -module,

$$\mathcal{S}(\mathcal{M}_{k,n}^*) = \mathbb{C}[x_{11}, \dots, x_{k1}] \otimes \cdots \otimes \mathbb{C}[x_{1n}, \dots, x_{kn}] \quad (3.6)$$

¹I am grateful to M. Vergne for pointing out that the decomposition (3.4) was known long before [Zh] and the work of Howe [Ho] and to M. Brion for providing the reference [Mc].

and is therefore acted upon by the $\mathfrak{gl}_k$ -intertwiners $\tilde{\Omega}_{ij}^{(k)}$, $1 \leq i < j \leq n$, defined by

$$\tilde{\Omega}_{ij}^{(k)} = \sum_a 1^{\otimes(i-1)} \otimes X_a \otimes 1^{\otimes(j-i-1)} \otimes X^a \otimes 1^{\otimes(n-j)} \quad (3.7)$$

where $\{X_a\}$, $\{X^a\}$ are dual basis of $\mathfrak{gl}_k$ with respect to the pairing $\langle X, Y \rangle = \text{tr}(XY)$. On the other hand, as a $\mathfrak{gl}_n$ -module, $\mathcal{S}(\mathcal{M}_{k,n}^*)$ is acted upon by the operators $\kappa_{ij}^{(n)}$, $1 \leq i < j \leq n$, where

$$\kappa_{ij}^{(n)} = e_\alpha f_\alpha + f_\alpha e_\alpha \quad (3.8)$$

is the truncated Casimir operator of the $\mathfrak{sl}_2(\mathbb{C})$ -subalgebra of $\mathfrak{gl}_n$ corresponding to the root $\alpha = \theta_i - \theta_j$. Let $e_1, \dots, e_p$ be the canonical basis of $\mathbb{C}^p$ and $E_{ab}^{(p)} e_c = \delta_{bc} e_a$, $1 \leq a, b \leq p$ the corresponding basis of $\mathfrak{gl}_p$ with dual basis $E_{ba}^{(p)}$. Let $1 \leq i < j \leq n$, then

Proposition 3.3. *The following holds on $\mathcal{S}(\mathcal{M}_{k,n}^*)$*

$$2\tilde{\Omega}_{ij}^{(k)} = \kappa_{ij}^{(n)} - E_{ii}^{(n)} - E_{jj}^{(n)} \quad (3.9)$$

PROOF. By (3.7), $\tilde{\Omega}_{ij}^{(k)}$ acts on $\mathcal{S}(\mathcal{M}_{k,n}^*)$ as

$$\tilde{\Omega}_{ij}^{(k)} = \sum_{1 \leq a, b \leq k} x_{ai} \partial_{bi} x_{bj} \partial_{aj} \quad (3.10)$$

where x_{rc} and ∂_{rc} are the operators of multiplication by and derivation with respect to x_{rc} . On the other hand, given that the $\mathfrak{sl}_2(\mathbb{C})$ -triple $\{e_\alpha, h_\alpha, f_\alpha\}$ corresponding to the root $\alpha = \theta_i - \theta_j$ of $\mathfrak{gl}_n$ is $\{E_{ij}^{(n)}, E_{ii}^{(n)} - E_{jj}^{(n)}, E_{ji}^{(n)}\}$, the following holds on $\mathcal{S}(\mathcal{M}_{k,n}^*)$

$$\kappa_{ij}^{(n)} = \sum_{1 \leq a, b \leq k} x_{ai} \partial_{aj} x_{bj} \partial_{bi} + x_{bj} \partial_{bi} x_{ai} \partial_{aj} \quad (3.11)$$

Subtracting, we find

$$2\tilde{\Omega}_{ij}^{(k)} - \kappa_{ij}^{(n)} = - \sum_{1 \leq a, b \leq k} \delta_{ab} x_{ai} \partial_{bi} + \delta_{ab} x_{bj} \partial_{aj} = -E_{ii}^{(n)} - E_{jj}^{(n)} \quad (3.12)$$

as claimed ■

Let $\lambda \in \mathbb{Y}_{\min(k,n)}$ and $V_\lambda^{(n)}$ the corresponding simple GL_n -module. By theorem 3.2, $V_\lambda^{(n)}$ may be identified with the subspace of vectors of highest weight λ for the action of $\mathfrak{gl}_k$ on $\mathcal{S}(\mathcal{M}_{k,n}^*)$. Denote by $\iota : V_\lambda \rightarrow \mathcal{S}(\mathcal{M}_{k,n}^*)$ the corresponding $\mathfrak{gl}_n$ -equivariant embedding and let $\mu = (\mu_1, \dots, \mu_n) \in \mathbb{N}^n$ be a weight of $V_\lambda^{(n)}$.

Lemma 3.4. *The embedding ι maps the subspace $V_\lambda^{(n)}[\mu] \subset V_\lambda^{(n)}$ of weight μ onto the subspace M_λ^μ of vectors of highest weight λ for the action of $\mathfrak{gl}_k$ on*

$$S^\mu \mathbb{C}^k = S^{\mu_1} \mathbb{C}^k \otimes \dots \otimes S^{\mu_n} \mathbb{C}^k \subset \mathbb{C}[x_{11}, \dots, x_{k1}] \otimes \dots \otimes \mathbb{C}[x_{1n}, \dots, x_{kn}] \quad (3.13)$$

where $S^{\mu_j} \mathbb{C}^k$ is the space of polynomials in $x_{1j}, \dots, x_{kj}$ which are homogeneous of degree μ_j . The corresponding isomorphism

$$\bigoplus_{\nu \in \mathfrak{S}_n \mu} V_\lambda^{(n)}[\nu] \cong \bigoplus_{\nu \in \mathfrak{S}_n \mu} M_\lambda^\nu \quad (3.14)$$

is equivariant with respect to $\mathfrak{S}_n$ which acts on $\bigoplus_{\nu \in \mathfrak{S}_n \mu} S^\nu \mathbb{C}^k$ by permuting the tensor factors and on V_λ by regarding $\mathfrak{S}_n$ as the subgroup of permutation matrices of $GL_n(\mathbb{C})$.

PROOF. The equality $\iota(V_\lambda^{(n)}[\mu]) = M_\lambda^\mu$ holds because $S^\mu \mathbb{C}^k$ is the subspace of $\mathcal{S}(\mathcal{M}_{k,n}^*)$ of weight μ for the $\mathfrak{gl}_n$ -action since $E_{ii}^{(n)} x_{rj}^m = \delta_{ij} m x_{rj}^m$. The $\mathfrak{S}_n$ -equivariance stems from the fact that the permutation of the tensor factors in $S^\bullet \mathbb{C}^k \otimes \cdots \otimes S^\bullet \mathbb{C}^k \cong S^\bullet(\mathbb{C}^k \otimes \mathbb{C}^n)$ is given by the action of $\mathfrak{S}_n \subset GL_n(\mathbb{C})$ action on $\mathbb{C}^n$ ■

Let $\mathcal{D}_n = \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid z_i = z_j \text{ for some } 1 \leq i < j \leq n\}$ and $X_n = \mathbb{C}^n \setminus \mathcal{D}_n$. Regard $\mathbb{C}_0^n = \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid \sum_{j=1}^n z_j = 0\}$ as the Cartan subalgebra of diagonal matrices in $\mathfrak{sl}_n$ and $X_n^0 = \mathbb{C}_0^n \setminus \mathcal{D}_n$ as the set of its regular elements. Since the inclusion $X_n^0 \subset X_n$ is a homotopy equivalence, $\pi_1(X_n) \cong \pi_1(X_n^0) = B_n$ are generated by $T_1 \dots T_{n-1}$ with

$$T_i T_j = T_j T_i \quad \text{if } |i - j| \geq 2 \quad (3.15)$$

$$T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1} \quad i = 1 \dots n-1 \quad (3.16)$$

Define $\Omega_{ij}^{(k)} \in \text{End}_{\mathfrak{gl}_k}(S^\mu \mathbb{C}^k)$ by (3.7) where now $\{X_a\}, \{X^a\}$ are dual basis of $\mathfrak{sl}_k$ and extend the connection (2.4) to X_n in the obvious way. The following is the main result of this section.

Theorem 3.5. *$f : X_n \rightarrow M_\lambda^\mu \subset S^{\mu_1} \mathbb{C}^k \otimes \cdots \otimes S^{\mu_n} \mathbb{C}^k$ is a horizontal section of the Knizhnik–Zamolodchikov connection*

$$\nabla_{\text{KZ}}^{\bar{h}} = d - \bar{h} \sum_{1 \leq i < j \leq n} \frac{dz_i - dz_j}{z_i - z_j} \Omega_{ij}^{(k)} \quad (3.17)$$

iff the $V_\lambda^{(n)}[\mu]$ -valued function $g = f \cdot \prod_{1 \leq i < j \leq n} (z_i - z_j)^{h(\mu_i + \mu_j + 2\mu_i \mu_j / k)}$ is a horizontal section of

$$\nabla_\kappa^h = d - h \sum_{1 \leq i < j \leq n} \frac{dz_i - dz_j}{z_i - z_j} \kappa_{ij}^{(n)} \quad (3.18)$$

where $\bar{h} = 2h$.

PROOF. Let $1^{(k)} = \sum_{i=1}^k E_{ii}^{(k)}$ be the generator of the centre of $\mathfrak{gl}_k$ so that, in obvious notation, $\tilde{\Omega}_{ij}^{(k)} = \Omega_{ij}^{(k)} + \frac{1}{k} \pi_i(1^{(k)}) \pi_j(1^{(k)})$. The operators $2\tilde{\Omega}_{ij}^{(k)}$ and $\kappa_{ij}^{(n)}$ both act on $M_\lambda^\mu \cong V_\lambda^{(n)}[\mu]$ and, by proposition 3.3, their restrictions differ by $-\mu_i - \mu_j$. The claim follows since, for any $1 \leq l \leq n$, $\pi_l(1^{(k)})$ acts on $S^\mu \mathbb{C}^k$ as multiplication by μ_l ■

REMARK. When $k = 2$ and λ is of the form $(|\mu|/2, |\mu|/2, 0, \dots, 0)$, where $|\mu| = \sum_{i=1} \mu_i$, theorem 3.5 is a representation-theoretic analogue of the coincidence between the Kapovich–Millson bending flows on the space of n -gons in $\mathbb{R}^3$ with side lengths $\mu_1, \dots, \mu_n$ [KM] and the Gel'fand–Cetlin flows on the Grassmannian $\text{Gr}_2(\mathbb{C}^n)$ [GS] observed by Hausmann and Knutson in the context of Gel'fand–McPherson duality [HK]. I am grateful to J. Millson for a careful explanation of this coincidence.

REMARK. An interesting relation between ∇_κ and the Knizhnik–Zamolodchikov connection was recently noted by Felder, Markov, Tarasov and Varchenko in [FMTV], where a variant of the connection (2.4) is independently introduced and studied. One of the main results of [FMTV] is that, for any simple Lie algebra $\mathfrak{g}$, the connection ∇_κ with values in a tensor product $V_1 \otimes \cdots \otimes V_n$ of n simple $\mathfrak{g}$ -modules is, when supplemented by suitable dynamical parameters, bispectral to (i.e., commutes with) the Knizhnik–Zamolodchikov connection for $\mathfrak{g}$ with values in the same n -fold tensor product. An analogous result is obtained in [TV] for a difference analogue of the connection ∇_κ . By comparison, theorem 3.5 can only hold for $\mathfrak{g} = \mathfrak{sl}_n$, since it relies on the 'coincidence' of the regular Cartan of $\mathfrak{gl}_n$ with the configuration

space of n ordered points in $\mathbb{C}$, and asserts the *equality* of the two connections.

To relate the monodromy representations of B_n corresponding to $\nabla_{\text{KZ}}^{\bar{h}}$ and ∇_{κ}^h , we need to specify how these induce flat connections on $X_n/\mathfrak{S}_n$ and $X_n^0/\mathfrak{S}_n$ respectively. For $\nabla_{\text{KZ}}^{\bar{h}}$, we let $\mathfrak{S}_n$ act on the fibre

$$\bigoplus_{\nu \in \mathfrak{S}_n \mu} M_{\lambda}^{\nu} \subset \bigoplus_{\nu \in \mathfrak{S}_n \mu} \mathcal{S}^{\nu} \mathbb{C}^k \quad (3.19)$$

by permuting the tensor factors and take the quotient connection. For ∇_{κ}^h , we use the construction of proposition 2.3 and the homomorphism $\sigma : B_n \rightarrow SL_n(\mathbb{C})$ given by

$$T_j \longrightarrow \exp(E_{j,j+1}^{(n)}) \exp(-E_{j+1,j}^{(n)}) \exp(E_{j,j+1}^{(n)}) = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & 0 & 1 & \\ & & & -1 & 0 & \\ & & & & & 1 & \\ & & & & & & \ddots & \\ & & & & & & & 1 \end{pmatrix} \quad (3.20)$$

where the off-diagonal terms are the $(j, j+1)$ and $(j+1, j)$ entries. A direct computation, or [Ti, thm. 3.3], show that the assignment (3.20) does indeed extend to a homomorphism $B_n \rightarrow SL_n(\mathbb{C})$. Choose the generators $T_1 \dots T_{n-1}$ of B_n as at the end of section 2.

Corollary 3.6. *Let μ be a weight of V_{λ} and*

$$\pi_{\kappa}^h : B_n \rightarrow GL\left(\bigoplus_{\nu \in \mathfrak{S}_n \mu} V_{\lambda}[\nu]\right), \quad \pi_{\text{KZ}}^{\bar{h}} : B_n \rightarrow GL\left(\bigoplus_{\nu \in \mathfrak{S}_n \mu} M_{\lambda}^{\nu}\right) \quad (3.21)$$

the monodromy representations of the braid group B_n corresponding to the connections (3.18) and (3.17) respectively. Then, for any $j = 1 \dots n-1$,

$$\pi_{\text{KZ}}^{2h}(T_j) = \pi_{\kappa}^h(T_j) \cdot e^{-\pi i h(E_{jj}^{(n)} + E_{j+1,j+1}^{(n)} + 2E_{jj}^{(n)} E_{j+1,j+1}^{(n)}/k)} \cdot e^{i\pi E_{jj}^{(n)}} \quad (3.22)$$

PROOF. Let $s_j \in SL_n(\mathbb{C})$ be the right-hand side of (3.20) so that $s_j = (j \ j+1) \cdot e^{i\pi E_{jj}^{(n)}}$ in $GL_n(\mathbb{C})$. Let $\mathcal{P}_{\text{KZ}}^{\bar{h}}, \mathcal{P}_{\kappa}^h$ denote parallel transport for $\nabla_{\text{KZ}}^{\bar{h}}$ and ∇_{κ}^h respectively. Then, by theorem 3.5 and proposition 2.4, the following holds on $M_{\lambda}^{\nu} \cong V_{\lambda}[\nu]$,

$$\begin{aligned} \pi_{\text{KZ}}^{2h}(T_j) &= (j \ j+1) \mathcal{P}_{\text{KZ}}^{2h}(T_j) \\ &= (j \ j+1) e^{-\pi i h(\nu_j + \nu_{j+1} + 2\nu_j \nu_{j+1}/k)} \mathcal{P}_{\kappa}^h(T_j) \\ &= s_j \mathcal{P}_{\kappa}^h(T_j) e^{i\pi E_{jj}^{(n)}} e^{-\pi i h(E_{jj}^{(n)} + E_{j+1,j+1}^{(n)} + 2E_{jj}^{(n)} E_{j+1,j+1}^{(n)}/k)} \\ &= \pi_{\kappa}^h(T_j) e^{-\pi i h(E_{jj}^{(n)} + E_{j+1,j+1}^{(n)} + 2E_{jj}^{(n)} E_{j+1,j+1}^{(n)}/k)} e^{i\pi E_{jj}^{(n)}} \end{aligned} \quad (3.23)$$

as claimed ■

4. THE QUANTUM GROUP $U_{\hbar} \mathfrak{gl}_p$

In this, and the following sections, we work over the ring $\mathbb{C}[[\hbar]]$ of formal power series in the variable $\hbar$. All tensor products of $\mathbb{C}[[\hbar]]$ -modules are understood to be completed in the $\hbar$ -adic topology. For $p \in \mathbb{N}$, let $a_{ij} = 2\delta_{ij} - \delta_{|i-j|=1}$, $1 \leq i, j \leq p$, be the entries of the Cartan matrix

of type A_{p-1} and let $U_{\hbar}\mathfrak{gl}_p$ be the corresponding Drinfeld–Jimbo quantum group [Dr1, Ji1] i.e., the algebra over $\mathbb{C}[[\hbar]]$ topologically generated by elements $E_i, F_i, i = 1 \dots p-1$ and $D_i, i = 1 \dots p$ subject to the q –Serre relations

$$[D_i, D_j] = 0 \quad (4.1)$$

$$[D_i, E_j] = (\delta_{ij} - \delta_{ij+1})E_j \quad [D_i, F_j] = -(\delta_{ij} - \delta_{ij+1})F_j \quad (4.2)$$

$$[E_i, F_j] = \delta_{ij} \frac{e^{\hbar H_i} - e^{-\hbar H_i}}{e^{\hbar} - e^{-\hbar}} \quad (4.3)$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix} E_i^k E_j E_i^{1-a_{ij}-k} = 0, \quad \forall i \neq j \quad (4.4)$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix} F_i^k F_j F_i^{1-a_{ij}-k} = 0, \quad \forall i \neq j \quad (4.5)$$

where $H_i = D_i - D_{i+1}$ and, for any $n \geq k \in \mathbb{N}$,

$$[n] = \frac{e^{n\hbar} - e^{-n\hbar}}{e^{\hbar} - e^{-\hbar}} \quad (4.6)$$

$$[n]! = [n][n-1] \cdots [1] \quad (4.7)$$

$$\begin{bmatrix} n \\ k \end{bmatrix} = \frac{[n]!}{[k]![n-k]!} \quad (4.8)$$

$U_{\hbar}\mathfrak{gl}_p$ is a topological Hopf algebra with coproduct Δ and counit ε given by

$$\Delta(D_i) = D_i \otimes 1 + 1 \otimes D_i \quad (4.9)$$

$$\Delta(E_i) = E_i \otimes e^{\hbar H_i} + 1 \otimes E_i \quad (4.10)$$

$$\Delta(F_i) = F_i \otimes 1 + e^{-\hbar H_i} \otimes F_i \quad (4.11)$$

and

$$\varepsilon(E_i) = \varepsilon(F_i) = \varepsilon(D_i) = 0 \quad (4.12)$$

Note that $I = D_1 + \cdots + D_p$ is central so that $U_{\hbar}\mathfrak{gl}_p \cong U_{\hbar}\mathfrak{sl}_p \otimes \mathbb{C}[I][[\hbar]]$ as Hopf algebras where the coproduct on $\mathbb{C}[I][[\hbar]]$ is given by $\Delta(I) = I \otimes 1 + 1 \otimes I$ and $U_{\hbar}\mathfrak{sl}_p \subset U_{\hbar}\mathfrak{gl}_p$ is the closed Hopf subalgebra generated by E_i, F_i and $H_i, i = 1 \dots p-1$.

By a finite–dimensional representation of $U_{\hbar}\mathfrak{gl}_p$ we shall mean a $U_{\hbar}\mathfrak{gl}_p$ –module which is topologically free and finitely generated over $\mathbb{C}[[\hbar]]$ and on which I acts semisimply with eigenvalues in $\mathbb{C}$. Choose an algebra isomorphism $\phi : U_{\hbar}\mathfrak{gl}_p \rightarrow U_{\hbar}\mathfrak{gl}_p[[\hbar]]$ mapping each D_i onto E_{ii} [Dr2, prop. 4.3] and let V be a finite–dimensional $\mathfrak{gl}_p$ –module on which $1^{(p)} = \sum_{i=1}^p E_{ii}$ acts semisimply. Then, $U_{\hbar}\mathfrak{gl}_p[[\hbar]]$ acts on $V[[\hbar]]$ and the latter becomes, via ϕ , a finite–dimensional representation of $U_{\hbar}\mathfrak{gl}_p$. Conversely,

Proposition 4.1. *Let $\mathcal{V}$ be a finite–dimensional representation of $U_{\hbar}\mathfrak{gl}_p$ and $V = \mathcal{V}/\hbar V$ the corresponding $\mathfrak{gl}_p$ –module. Then, as $U_{\hbar}\mathfrak{gl}_p$ –modules,*

$$\mathcal{V} \cong V[[\hbar]] \quad (4.13)$$

PROOF. Since I is diagonalisable on $\mathcal{V}$ and commutes with $U_{\hbar}\mathfrak{gl}_p$, we may assume that it acts on $\mathcal{V}$ as multiplication by a scalar $\lambda \in \mathbb{C}$. Since $\mathcal{V}$ is topologically free, $\mathcal{V} \cong V[[\hbar]]$ as $\mathbb{C}[[\hbar]]$ –modules so that $\mathcal{V}$ is a deformation of the finite–dimensional $\mathfrak{sl}_p$ –module V . Since $\mathfrak{sl}_p$ is

simple, $H^1(\mathfrak{sl}_p, V) = 0$ and $\mathcal{V}$ is isomorphic, as $\mathfrak{sl}_p$, and therefore as $\mathfrak{gl}_p$ -module to the trivial deformation of V . Thus, $\mathcal{V} \cong V[[\hbar]]$ as $U\mathfrak{gl}_p[[\hbar]]$, and therefore as $U_\hbar\mathfrak{gl}_p$ -modules ■

Corollary 4.2. *Let U, V be finite-dimensional $\mathfrak{gl}_p$ -modules on which $1^{(p)}$ acts semisimply. If $U \otimes V$ decomposes as*

$$U \otimes V \cong \bigoplus_W N_W W \quad (4.14)$$

for some $\mathfrak{gl}_p$ -modules W and multiplicities $N_W \in \mathbb{N}$, then, as $U_\hbar\mathfrak{gl}_p$ -modules,

$$U[[\hbar]] \otimes V[[\hbar]] \cong \bigoplus_W N_W W[[\hbar]] \quad (4.15)$$

PROOF. By (4.14), both sides of (4.15) have the same specialisation at $\hbar = 0$ and are therefore isomorphic by proposition 4.1 ■

5. THE DUAL PAIR $(U_\hbar\mathfrak{gl}_k, U_\hbar\mathfrak{gl}_n)$

We shall need the analogue of theorem 3.2 in the setting of the algebra $\mathcal{S}_\hbar(\mathcal{M}_{k,n}^*)$ of functions on quantum $k \times n$ matrix space. With the exception of theorems 5.4 and 5.5, this section follows [Ba, §1.5] (see also [Ga]). By definition, $\mathcal{S}_\hbar(\mathcal{M}_{k,n}^*)$ is the algebra over $\mathbb{C}[[\hbar]]$ topologically generated by elements X_{ij} , $1 \leq i \leq k$, $1 \leq j \leq n$ with relations

$$X_{ij}X_{kl} = \begin{cases} X_{kl}X_{ij} & \text{if } k > i \text{ and } l < j \text{ or } k < i \text{ and } l > j \\ e^{-\hbar}X_{kl}X_{ij} & \text{if } k > i \text{ and } l = j \text{ or } k = i \text{ and } l > j \\ X_{kl}X_{ij} - (e^\hbar - e^{-\hbar})X_{kj}X_{il} & \text{if } k > i \text{ and } l > j \end{cases} \quad (5.1)$$

$\mathcal{S}_\hbar(\mathcal{M}_{k,n}^*)$ is $\mathbb{N}$ -graded by decreeing that each X_{ij} is of degree 1 and we denote its homogeneous components by $\mathcal{S}_\hbar^d(\mathcal{M}_{k,n}^*)$, $d \in \mathbb{N}$. For any $k \times n$ matrix m with entries $m_{ij} \in \mathbb{N}$, set

$$\begin{aligned} \mathbf{X}^m &= X_{11}^{m_{11}} \dots X_{k1}^{m_{k1}} \dots X_{1n}^{m_{1n}} \dots X_{kn}^{m_{kn}} \\ &= X_{11}^{m_{11}} \dots X_{1n}^{m_{1n}} \dots X_{k1}^{m_{k1}} \dots X_{kn}^{m_{kn}} \end{aligned} \quad (5.2)$$

By the commutation relations (5.1), the $\mathbf{X}^m$, with $m \in \mathcal{M}_{k,n}$ such that $|m| = d$, span $\mathcal{S}_\hbar^d(\mathcal{M}_{k,n}^*)$, where $|m| = \sum_{i,j} m_{ij}$.

Theorem 5.1 (Parshall–Wang). *The monomials $\mathbf{X}^m$, $m \in \mathcal{M}_{kn}(\mathbb{N})$, are linearly independent over $\mathbb{C}[[\hbar]]$. In particular, the set $\{\mathbf{X}^m\}_{|m|=d}$ is a $\mathbb{C}[[\hbar]]$ -basis of $\mathcal{S}_\hbar^d(\mathcal{M}_{k,n}^*)$.*

PROOF. This is proved in [PW, thm. 3.5.1] for $k = n$ and over the field $\mathbb{C}(q)$ of rational functions of $q = e^\hbar$ rather than over $\mathbb{C}[[\hbar]]$. The proof however works equally well for $k \neq n$ and, as remarked in [PW], over $\mathbb{C}[[\hbar]]$ ■

As in the classical case, $\mathcal{S}_\hbar(\mathcal{M}_{k,n}^*)$ is a module algebra over $U_\hbar\mathfrak{gl}_k \otimes U_\hbar\mathfrak{gl}_n$. This may be seen in the following way. For any $l \in \mathbb{N}$, one readily checks that the assignement

$$X_{ij} \rightarrow \sum_{l'=1}^l X_{il'} \otimes X_{l'j} \quad (5.3)$$

extends uniquely to an algebra homomorphism $\Delta_{kln} : \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \rightarrow \mathcal{S}_{\hbar}(\mathcal{M}_{k,l}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{l,n}^*)$ such that, for any $l, m \in \mathbb{N}$, the following diagram commutes

$$\begin{array}{ccc}
 \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) & \xrightarrow{\Delta_{kln}} & \mathcal{S}_{\hbar}(\mathcal{M}_{k,l}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{l,n}^*) \\
 \Delta_{kmn} \downarrow & & \downarrow 1 \otimes \Delta_{lmn} \\
 \mathcal{S}_{\hbar}(\mathcal{M}_{k,m}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{m,n}^*) & \xrightarrow{\Delta_{klm} \otimes 1} & \mathcal{S}_{\hbar}(\mathcal{M}_{k,l}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{l,m}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{m,n}^*)
 \end{array} \tag{5.4}$$

In particular, $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ and $\mathcal{S}_{\hbar}(\mathcal{M}_{n,n}^*)$ are topological bialgebras with comultiplications Δ_{kkk} and Δ_{nnn} respectively and counit $\varepsilon(X_{ij}) = \delta_{ij}$. Moreover, the maps Δ_{kkn} and Δ_{knn} give $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ the structure of a $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ - $\mathcal{S}_{\hbar}(\mathcal{M}_{n,n}^*)$ bicomodule algebra each homogeneous component of which is invariant under $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ and $\mathcal{S}_{\hbar}(\mathcal{M}_{n,n}^*)$ since $\Delta_{kln}(\mathcal{S}_{\hbar}^d(\mathcal{M}_{k,k}^*)) \subset \mathcal{S}_{\hbar}^d(\mathcal{M}_{k,k}^*) \otimes \mathcal{S}_{\hbar}^d(\mathcal{M}_{l,l}^*)$.

We shall need a columnwise (resp. rowwise) description of the coaction of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ (resp. $\mathcal{S}_{\hbar}(\mathcal{M}_{n,n}^*)$) on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$. Consider the quantum k and n -dimensional planes *i.e.*, the algebras $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)$ and $\mathcal{S}_{\hbar}(\mathcal{M}_{1,n}^*)$. By the commutation relations (5.1) and theorem 5.1, these may be embedded as subalgebras of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ via the maps

$$c_j : \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*) \rightarrow \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*), \quad c_j(X_{i1}) = X_{ij} \tag{5.5}$$

$$r_i : \mathcal{S}_{\hbar}(\mathcal{M}_{1,n}^*) \rightarrow \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*), \quad r_i(X_{1j}) = X_{ij} \tag{5.6}$$

with $1 \leq i \leq k, 1 \leq j \leq n$. By (5.4), $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)$ is a left algebra comodule over $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ and $\mathcal{S}_{\hbar}(\mathcal{M}_{1,n}^*)$ a right algebra comodule over $\mathcal{S}_{\hbar}(\mathcal{M}_{n,n}^*)$.

Lemma 5.2. *As left, $\mathbb{N}$ -graded $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ -comodules,*

$$\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \cong \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n} \tag{5.7}$$

via the map $\Phi : p_1 \otimes \cdots \otimes p_n \rightarrow c_1(p_1) \cdots c_n(p_n)$. Similarly, as right, $\mathbb{N}$ -graded $\mathcal{S}_{\hbar}(\mathcal{M}_{n,n}^*)$ -comodules,

$$\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \cong \mathcal{S}_{\hbar}(\mathcal{M}_{1,n}^*)^{\otimes k} \tag{5.8}$$

via $\Psi : q_1 \otimes \cdots \otimes q_k \rightarrow r_1(q_1) \cdots r_k(q_k)$.

PROOF. The map Φ clearly preserves the grading and, by theorem 5.1, restricts to a $\mathbb{C}[[\hbar]]$ -linear isomorphism of homogeneous components since it bijectively maps the monomial basis of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n}$ onto the basis $\mathbf{X}^m$ of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$. The fact that Φ itself is an isomorphism follows easily because any element of $\mathcal{S}_{\hbar}(\mathcal{M}_{1,n}^*)^{\otimes k}$ or $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ is the convergent sum of its homogeneous components. As readily checked, the diagram

$$\begin{array}{ccc}
 \mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*) & \xrightarrow{1 \otimes c_j} & \mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \\
 \Delta_{kk1} \uparrow & & \uparrow \Delta_{kkn} \\
 \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*) & \xrightarrow{c_j} & \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)
 \end{array} \tag{5.9}$$

is commutative for any $1 \leq j \leq n$ and therefore so is

$$\begin{array}{ccc}
 (\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*))^{\otimes n} & \xrightarrow{\mu^{(n)} \otimes \Phi} & \mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \\
 \Delta_{kk1}^{\otimes n} \uparrow & & \uparrow \Delta_{kkn} \\
 \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n} & \xrightarrow{\Phi} & \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)
 \end{array} \tag{5.10}$$

where $\mu^{(n)} : \mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)^{\otimes n} \rightarrow \mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ is the n -fold multiplication. This proves (5.7). The proof of (5.8) is identical ■

We turn now to the action of $U_{\hbar}\mathfrak{gl}_k \otimes U_{\hbar}\mathfrak{gl}_n$ on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$. For $p = k, n$, consider the vector representation of $U_{\hbar}\mathfrak{gl}_p$ i.e., the module $V = \mathbb{C}^p[[\hbar]]$ with basis $e_1, \dots, e_p$ and action given by

$$D_i = E_{ii}, \quad E_i = E_{ii+1}, \quad F_i = E_{i+1i} \tag{5.11}$$

where $E_{ab}e_c = \delta_{bc}e_a$. Let $e^1, \dots, e^p \in V^*$ be the dual basis of $e_1, \dots, e_p$, $U_{\hbar}\mathfrak{gl}_p^{\circ}$ the restricted dual of $U_{\hbar}\mathfrak{gl}_p$ and $t_{ij} \in U_{\hbar}\mathfrak{gl}_p^{\circ}$ the matrix coefficient defined by

$$t_{ij}(x) = \langle e^i, x e_j \rangle \tag{5.12}$$

Proposition 5.3. *The assignment $X_{ij} \rightarrow t_{ij}$ extends uniquely to a bialgebra morphism $\kappa_p : \mathcal{S}_{\hbar}(\mathcal{M}_{p,p}^*) \rightarrow U_{\hbar}\mathfrak{gl}_p^{\circ}$.*

PROOF. We need to check that the t_{ij} satisfy the relations (5.1), i.e., that when evaluated on $\Delta(x)$, $x \in U_{\hbar}\mathfrak{gl}_p$,

$$t_{ij} \otimes t_{kl} = \begin{cases} t_{kl} \otimes t_{ij} & \text{if } k > i \text{ and } l < j \text{ or } k < i \text{ and } l > j \\ e^{-\hbar} t_{kl} \otimes t_{ij} & \text{if } k > i \text{ and } l = j \text{ or } k = i \text{ and } l > j \\ t_{kl} \otimes t_{ij} - (e^{\hbar} - e^{-\hbar}) t_{kj} \otimes t_{il} & \text{if } k > i \text{ and } l > j \end{cases} \tag{5.13}$$

Let $R^{\vee} = \sigma \cdot R \in \text{End}(V \otimes V)$ where $\sigma \in GL(V \otimes V)$ is the flip and R is the universal R -matrix of $U_{\hbar}\mathfrak{gl}_p$ acting on $V \otimes V$. Then [Ji2],[CP, §8.3.G]

$$R = \left(e^{\hbar} \sum_{i=1}^p E_{ii} \otimes E_{ii} + \sum_{1 \leq i \neq j \leq p} E_{ii} \otimes E_{jj} + (e^{\hbar} - e^{-\hbar}) \sum_{1 \leq i < j \leq p} E_{ij} \otimes E_{ji} \right) \tag{5.14}$$

so that the matrix entries of $R^{\vee}$ are

$$R_{ik,jl}^{\vee} = \begin{cases} e^{\delta_{jl}\hbar} & \text{if } i = l \text{ and } k = j \\ e^{\hbar} - e^{-\hbar} & \text{if } i = j, k = l \text{ and } j > l \\ 0 & \text{otherwise} \end{cases} \tag{5.15}$$

From (5.15), one readily checks that both sides of (5.13) coincide when evaluated on any $A \in \text{End}(V \otimes V)$ commuting with $R^{\vee}$. Since $R^{\vee}$ is a $U_{\hbar}\mathfrak{gl}_p$ -intertwiner, κ extends to an algebra morphism which respects the counit and coproduct since $\Delta(t_{ij}) = \sum_{q=1}^p t_{iq} \otimes t_{qj}$ ■

Theorem 5.4.

1. *The maps κ_p , $p = k, n$ of proposition 5.3 give $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ the structure of an algebra module over $U_{\hbar}\mathfrak{gl}_k \otimes U_{\hbar}\mathfrak{gl}_n$ with invariant homogeneous components $\mathcal{S}_{\hbar}^d(\mathcal{M}_{k,n}^*)$, $d \in \mathbb{N}$.*

2. The maps Φ, Ψ of lemma 5.2 yield isomorphisms

$$\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \cong \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n} \quad \text{and} \quad \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \cong \mathcal{S}_{\hbar}(\mathcal{M}_{1,n}^*)^{\otimes k} \quad (5.16)$$

as $\mathbb{N}$ -graded $U_{\hbar}\mathfrak{gl}_k$ and $U_{\hbar}\mathfrak{gl}_n$ -modules respectively.

3. The action of the generators $E_q^{(p)}, F_q^{(p)}$, $q = 1 \dots p-1$ and $D_q^{(p)}$, $q = 1 \dots p$ of $U_{\hbar}\mathfrak{gl}_p$, $p = k, n$ in the monomial basis $\mathbf{X}^m$, $m \in \mathcal{M}_{k,n}$, is given by

$$D_i^{(k)} \mathbf{X}^m = \sum_{j=1}^n m_{ij} \mathbf{X}^m \quad (5.17)$$

$$E_i^{(k)} \mathbf{X}^m = \sum_{j=1}^n [m_{i+1j}] \prod_{j'=j+1}^n e^{\hbar(m_{ij'} - m_{i+1j'})} \mathbf{X}^{m + \varepsilon_{ij} - \varepsilon_{i+1j}} \quad (5.18)$$

$$F_i^{(k)} \mathbf{X}^m = \sum_{j=1}^n [m_{ij}] \prod_{j'=1}^{j-1} e^{-\hbar(m_{ij'} - m_{i+1j'})} \mathbf{X}^{m - \varepsilon_{ij} + \varepsilon_{i+1j}} \quad (5.19)$$

where $(\varepsilon_{ab})_{cd} = \delta_{ac}\delta_{bd}$, and

$$D_j^{(n)} \mathbf{X}^m = \sum_{i=1}^k m_{ij} \mathbf{X}^m \quad (5.20)$$

$$E_j^{(n)} \mathbf{X}^m = \sum_{i=1}^k [m_{ij+1}] \prod_{i'=i+1}^k e^{\hbar(m_{i'j} - m_{i'j+1})} \mathbf{X}^{m + \varepsilon_{ij} - \varepsilon_{ij+1}} \quad (5.21)$$

$$F_j^{(n)} \mathbf{X}^m = \sum_{i=1}^k [m_{ij}] \prod_{i'=1}^{i-1} e^{-\hbar(m_{i'j} - m_{i'j+1})} \mathbf{X}^{m - \varepsilon_{ij} + \varepsilon_{ij+1}} \quad (5.22)$$

PROOF. Using the transposition anti-involution τ on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*)$ given by $\tau(X_{ij}) = X_{ji}$, we may regard $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ as a right algebra comodule over $\mathcal{S}_{\hbar}(\mathcal{M}_{k,k}^*) \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{n,n}^*)$ and therefore, via the pairings $\langle \cdot, \cdot \rangle : \mathcal{S}_{\hbar}(\mathcal{M}_{m,m}^*) \otimes U_{\hbar}\mathfrak{gl}_m \rightarrow \mathbb{C}[[\hbar]]$, $m = k, n$, given by proposition 5.3 as a left algebra module over $U_{\hbar}\mathfrak{gl}_k \otimes U_{\hbar}\mathfrak{gl}_n$. This proves (i) and (ii). Explicitly, for $x^{(m)} \in U_{\hbar}\mathfrak{gl}_m$, $m = k, n$ and $p \in \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$

$$x^{(k)} p = \langle x^{(k)} \otimes 1, \tau \otimes 1 \cdot \Delta_{kkn}(p) \rangle \quad (5.23)$$

$$x^{(n)} p = \langle 1 \otimes x^{(n)}, \Delta_{knn}(p) \rangle \quad (5.24)$$

Using (5.23) and (5.11), one gets

$$D_i^{(k)} X_{i'j} = \delta_{ii'} X_{ij} \quad (5.25)$$

$$E_i^{(k)} X_{i'j} = \delta_{i+1i'} X_{ij} \quad (5.26)$$

$$F_i^{(k)} X_{i'j} = \delta_{ii'} X_{i+1j} \quad (5.27)$$

Using the algebra module property $x(pq) = \mu(\Delta(x)p \otimes q)$ where $x \in U_{\hbar}\mathfrak{gl}_k$, $p, q \in \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ and $\mu : \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)^{\otimes 2} \rightarrow \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ is multiplication, (4.9)–(4.11) and the commutation relations (5.1) shows by induction on $m \in \mathbb{N}$ that

$$D_i^{(k)} X_{i'j}^m = \delta_{ii'} m X_{ij}^m \quad (5.28)$$

$$E_i^{(k)} X_{i'j}^m = \delta_{i+1i'} [m] X_{ij} X_{i+1j}^{m-1} \quad (5.29)$$

$$F_i^{(k)} X_{i'j}^m = \delta_{ii'} [m] X_{ij}^{m-1} X_{i+1j} \quad (5.30)$$

Let $\Delta^{(a)} : U_{\hbar}\mathfrak{gl}_k \rightarrow U_{\hbar}\mathfrak{gl}_k^{\otimes a}$, $a \in \mathbb{N}^*$ be recursively defined by $\Delta^{(1)} = \text{id}$, $\Delta^{(a+1)} = \Delta \otimes \text{id}^{\otimes(a-1)} \cdot \Delta^{(a)}$. Then, by (4.9)–(4.11)

$$\Delta^{(a)} D_i^{(k)} = \sum_{b=1}^a 1^{\otimes(b-1)} \otimes D_i^{(k)} \otimes 1^{\otimes(a-b)} \quad (5.31)$$

$$\Delta^{(a)} E_i^{(k)} = \sum_{b=1}^a 1^{\otimes(b-1)} \otimes E_i^{(k)} \otimes (e^{\hbar H_i^{(k)}})^{\otimes(a-b)} \quad (5.32)$$

$$\Delta^{(a)} F_i^{(k)} = \sum_{b=1}^a (e^{-\hbar H_i^{(k)}})^{\otimes(b-1)} \otimes F_i^{(k)} \otimes 1^{\otimes(a-b)} \quad (5.33)$$

The formulae (5.17)–(5.19) now follow from the algebra module property and (5.31)–(5.33). The proof of (5.20)–(5.22) is similar ■

The following result is proved in [Ba] and [Ga] for the quantum groups $U_q\mathfrak{gl}_k, U_q\mathfrak{gl}_n$ by a different method

Theorem 5.5. *For any $d \in \mathbb{N}$, the $U_{\hbar}\mathfrak{gl}_k \otimes U_{\hbar}\mathfrak{gl}_n$ -module $\mathcal{S}_{\hbar}^d(\mathcal{M}_{k,n}^*)$ decomposes as*

$$\mathcal{S}_{\hbar}^d(\mathcal{M}_{k,n}^*) \cong \bigoplus_{\substack{\lambda \in \mathbb{Y}_{\min(k,n)} \\ |\lambda|=d}} V_{\lambda}^{(k)}[\hbar] \otimes V_{\lambda}^{(n)}[\hbar] \quad (5.34)$$

PROOF. By theorem 5.1, $\mathcal{S}_{\hbar}^d(\mathcal{M}_{k,n}^*)$ has no torsion, and is therefore a topologically free $\mathbb{C}[[\hbar]]$ -module. Moreover, by (5.17)–(5.22), $\mathcal{S}_{\hbar}^d(\mathcal{M}_{k,n}^*)/\hbar\mathcal{S}_{\hbar}^d(\mathcal{M}_{k,n}^*)$ is the $\mathfrak{gl}_k \oplus \mathfrak{gl}_n$ -module $\mathcal{S}^d(\mathcal{M}_{k,n}^*)$. The conclusion follows from theorem 3.2 and proposition 4.1 ■

6. BRAID GROUP ACTIONS ON QUANTUM MATRIX SPACE

We compare in this section two actions of the braid group B_n on the algebra $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ of functions of quantum $k \times n$ matrix space. The first is the R -matrix representation obtained by regarding $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ as the $U_{\hbar}\mathfrak{gl}_k$ -module $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n}$. The second is the quantum Weyl group action of B_n on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ viewed as a $U_{\hbar}\mathfrak{gl}_n$ -module. We will show that these representations essentially coincide, thus extending to the q -setting the fact that the symmetric group $\mathfrak{S}_n$ acts on $(S^{\bullet}\mathbb{C}^k)^{\otimes n} \cong \mathcal{S}(\mathcal{M}_{k,n}^*)$ via the permutation matrices in $GL_n(\mathbb{C})$.

More precisely, for any $1 \leq j \leq n$, let $R_j^{\vee}$ be the universal R -matrix of $U_{\hbar}\mathfrak{gl}_k$ acting on the j and $j+1$ tensor copies of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \cong \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n}$ and S_j the quantum Weyl group element of $U_{\hbar}\mathfrak{gl}_n$ corresponding to the simple root $\alpha_j = \theta_j - \theta_{j+1}$. We will show that

$$R_j^{\vee} = S_j \cdot e^{-\hbar(D_j^{(n)} + D_j^{(n)} D_{j+1}^{(n)}/k)} \cdot e^{i\pi D_j^{(n)}} \quad (6.1)$$

where $D_1^{(n)}, \dots, D_n^{(n)}$ are the generators of the Cartan subalgebra of $U_{\hbar}\mathfrak{gl}_n$. The proof of (6.1) is based upon the following observation, which we owe to B. Feigin. Both sides of (6.1) only act upon the j and $j+1$ tensor copies of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n}$ so that its proof reduces to a computation in $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes 2} \cong \mathcal{S}_{\hbar}(\mathcal{M}_{k,2}^*)$. Since both sides intertwine the action of $U_{\hbar}\mathfrak{gl}_k$ on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,2}^*)$, it suffices to compare them on highest weight vectors. These, and the action of $R_j^{\vee}$ are computed in §6.1. The action of S_j is computed in §6.2.

REMARK. It is easy to check that neither action of B_n is compatible with the algebra structure of $\mathcal{S}_h(\mathcal{M}_{k,n}^*)$, so that (6.1) cannot be proved by merely checking it on the generators X_{ij} . This stems from the fact that quantum Weyl group operators are not group-like.

6.1. R -matrix action on singular vectors. For any $d \in \mathbb{N}$, let $\mathcal{S}_h^d \mathbb{C}^k$ be the homogeneous component of degree d of $\mathcal{S}_h(\mathcal{M}_{k,1}^*)$. By (5.17)–(5.19), $\mathcal{S}_h^d \mathbb{C}^k$ is a deformation of the d -th symmetric power $\mathcal{S}^d \mathbb{C}^k$ of the vector representation of $\mathfrak{gl}_k$. Let $\mu_1, \mu_2 \in \mathbb{N}$, then

Lemma 6.1. *As $U_h \mathfrak{gl}_k$ -modules,*

$$S_h^{\mu_1} \mathbb{C}^k \otimes S_h^{\mu_2} \mathbb{C}^k \cong \bigoplus_{i=0}^{\min(\mu_1, \mu_2)} V_{\binom{\mu_1 + \mu_2 - i}{i}}^{(k)}[\hbar] \quad (6.2)$$

where $V_{\binom{\mu_1 + \mu_2 - i}{i}}^{(k)}$ is the irreducible representation of $\mathfrak{gl}_k$ with highest weight $(a, b, 0, \dots, 0)$. The corresponding highest weight vectors $v_i^{\mu_1, \mu_2}$ are given by

$$v_i^{\mu_1, \mu_2} = \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} X_{11}^{\mu_1 - i + a} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a} X_{22}^a \quad (6.3)$$

PROOF. The decomposition (6.2) follows from the Pieri rules for $\mathfrak{gl}_k$ and corollary 4.2. Fix $i \in \{0, \dots, \min(\mu_1, \mu_2)\}$. By (5.17), any $v \in S_h^{\mu_1} \mathbb{C}^k \otimes S_h^{\mu_2} \mathbb{C}^k$ of weight $(\mu_1 + \mu_2 - i, i, 0, \dots, 0)$ is of the form

$$v = \sum_{a=0}^i c_a X_{11}^{\mu_1 - i + a} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a} X_{22}^a \quad (6.4)$$

for some constants $c_a \in \mathbb{C}$. By (4.10) and (5.18), $\Delta(E_j)v = 0$ for any $j \geq 2$ so that v is a highest weight vector iff

$$\begin{aligned} \Delta(E_1)v &= \sum_{a=0}^i c_a [i - a] e^{\hbar(\mu_2 - 2a)} X_{11}^{\mu_1 - i + a + 1} X_{21}^{i - a - 1} \otimes X_{12}^{\mu_2 - a} X_{22}^a \\ &\quad + \sum_{a=0}^i c_a [a] X_{11}^{\mu_1 - i + a} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a + 1} X_{22}^{a - 1} \end{aligned} \quad (6.5)$$

is equal to zero. This yields $c_a = -c_{a-1} e^{\hbar(\mu_2 - 2a + 2)} [i - a + 1]/[a]$ and therefore

$$c_a = (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} c_0 \quad (6.6)$$

whence (6.3) ■

Let R be the universal R -matrix of $U_h \mathfrak{sl}_k$ and $R^\vee = \sigma \cdot R : S_h^{\mu_1} \mathbb{C}^k \otimes S_h^{\mu_2} \mathbb{C}^k \rightarrow S_h^{\mu_2} \mathbb{C}^k \otimes S_h^{\mu_1} \mathbb{C}^k$ the corresponding $U_h \mathfrak{sl}_k$ -intertwiner, where σ is the permutation of the tensor factors.

Proposition 6.2. *The following holds on $S_h^{\mu_1} \mathbb{C}^k \otimes S_h^{\mu_2} \mathbb{C}^k \oplus S_h^{\mu_2} \mathbb{C}^k \otimes S_h^{\mu_1} \mathbb{C}^k$,*

$$R^\vee v_i^{\mu_1, \mu_2} = (-1)^i e^{\hbar((\mu_1 - i)(\mu_2 - i) - i - \mu_1 \mu_2 / k)} v_i^{\mu_2, \mu_1} \quad (6.7)$$

PROOF. For any $1 \leq i \leq k - 1$, let $s_i = (i \ i + 1) \in \mathfrak{S}_k$ be the i th elementary transposition and let

$$w_0 = (1 \ k)(2 \ k - 1) \cdots (\lfloor \frac{k}{2} \rfloor \ \lceil \frac{k}{2} \rceil) \quad (6.8)$$

be the longest element of $\mathfrak{S}_k$. Consider the following reduced expression for w_0

$$w_0 = s_{k-1} \cdots s_1 s_{k-1} \cdots s_2 \cdots s_{k-1} s_{k-2} s_{k-1} = s_{i_1} \cdots s_{i_{k(k-1)/2}} \quad (6.9)$$

and let $\beta_j = s_{i_1} \cdots s_{i_{j-1}}(\theta_{i_j} - \theta_{i_{j+1}})$ be the associated enumeration of the positive roots of $\mathfrak{sl}_k$ so that

$$\begin{aligned} \beta_1 &= \theta_{k-1} - \theta_k, & \beta_2 &= \theta_{k-2} - \theta_k, & \cdots & & \beta_{k-1} &= \theta_1 - \theta_k, \\ & & \beta_k &= \theta_{k-2} - \theta_{k-1}, & \cdots & & \beta_{2k-3} &= \theta_1 - \theta_{k-1}, \\ & & & & \ddots & & & \\ & & & & & & \beta_{k(k-1)/2} &= \theta_1 - \theta_2 \end{aligned} \quad (6.10)$$

Let $E_{\beta_j}, F_{\beta_j} \in U_{\hbar} \mathfrak{sl}_k$, $1 \leq j \leq k(k-1)/2$ be the corresponding quantum root vectors so that $E_{\beta_j} = E_i$ and $F_{\beta_j} = F_i$ whenever β_j is the simple root α_i [Lu2, prop. 1.8]. Then, [KR, LS, Ro], [CP, thm. 8.3.9]

$$R = \exp \left(\hbar \sum_{i=1}^{k-1} H^i \otimes H_i \right) \prod_{j=1}^{k(k-1)/2} \exp_q((q - q^{-1})E_{\beta_j} \otimes F_{\beta_j}) \quad (6.11)$$

where $\{H^i\}_{i=1}^{k-1} \subset \mathfrak{h}$ is the basis of the Cartan subalgebra of $\mathfrak{sl}_k$ dual to $\{H_i\}_{i=1}^{k-1}$ with respect to the pairing $\langle X, Y \rangle = \text{tr}(XY)$, $q = e^{\hbar}$,

$$\exp_q(x) = \sum_{n \geq 0} q^{n(n-1)/2} \frac{x^n}{[n]!} \quad (6.12)$$

and the product in (6.11) is taken so that the factor $\exp_q((q - q^{-1})E_{\beta_j} \otimes F_{\beta_j})$ is placed to the left of $\exp_q((q - q^{-1})E_{\beta_{j'}} \otimes F_{\beta_{j'}})$ whenever $j > j'$. To compute $R v_i^{\mu_1, \mu_2}$, note that for any positive root $\beta \neq \theta_1 - \theta_2$ and $0 \leq a \leq \mu_1$,

$$E_{\beta} X_{11}^{\mu_1 - a} X_{21}^a = 0 \quad (6.13)$$

since, by (5.17), $(\mu_1 - a, a, 0, \dots, 0) + \beta$ is not a weight of $S_{\hbar}^{\mu_1} \mathbb{C}^k$. Thus, using (5.17)–(5.19)

$$\begin{aligned} R v_i^{\mu_1, \mu_2} &= \exp \left(\hbar \sum_{i=1}^{k-1} H^i \otimes H_i \right) \exp_q((q - q^{-1})E_1 \otimes F_1) v_i^{\mu_1, \mu_2} \\ &= \sum_{\substack{0 \leq a \leq i \\ 0 \leq n \leq i-a}} (-1)^a e^{\hbar a(\mu_2 - a + 1)} \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar((\mu_1 - i + a + n)(\mu_2 - a - n) + (i - a - n)(a + n) - \mu_1 \mu_2 / k)} \\ &\quad \cdot \frac{e^{\hbar n(n-1)/2} (e^{\hbar} - e^{-\hbar})^n}{[n]!} \frac{[i - a]! [\mu_2 - a]!}{[i - a - n]! [\mu_2 - a - n]!} \\ &\quad \cdot X_{11}^{\mu_1 - i + a + n} X_{21}^{i - a - n} \otimes X_{12}^{\mu_2 - a - n} X_{22}^{a + n} \end{aligned} \quad (6.14)$$

which, upon setting $\alpha = a + n$, yields

$$e^{\hbar((\mu_1 - i)(\mu_2 - i) - i - \mu_1 \mu_2 / k)} \sum_{\alpha=0}^i (-1)^{\alpha} \begin{bmatrix} i \\ \alpha \end{bmatrix} e^{\hbar(i - \alpha)(\mu_1 - i + \alpha + 1)} S_{\alpha}^{\mu_2} X_{11}^{\mu_1 - i + \alpha} X_{21}^{i - \alpha} \otimes X_{12}^{\mu_2 - \alpha} X_{22}^{\alpha} \quad (6.15)$$

where

$$S_{\alpha}^{\mu} = e^{\hbar \alpha(\mu - \alpha + 1)} \sum_{n=0}^{\alpha} (-1)^n \begin{bmatrix} \alpha \\ n \end{bmatrix} \frac{[\mu - \alpha + n]!}{[\mu - \alpha]!} e^{\hbar(\alpha - n)(\mu - \alpha + n + 1) + \hbar n(n-1)/2} (e^{\hbar} - e^{-\hbar})^n \quad (6.16)$$

We claim that $S_\alpha^\mu = 1$ for any $\alpha \leq \mu \in \mathbb{N}$ so that (6.7) holds. Indeed, using

$$\begin{bmatrix} \alpha \\ a \end{bmatrix} = \delta_{\alpha > a} \begin{bmatrix} \alpha - 1 \\ a \end{bmatrix} e^{-\hbar a} + \delta_{a > 0} \begin{bmatrix} \alpha - 1 \\ a - 1 \end{bmatrix} e^{\hbar(\alpha - a)} \quad (6.17)$$

one readily finds

$$S_\alpha^\mu = e^{2\hbar(\mu - \alpha + 1)} S_{\alpha - 1}^{\mu - 1} - (e^{2\hbar(\mu - \alpha + 1)} - 1) S_{\alpha - 1}^\mu \quad (6.18)$$

whence $S_\alpha^\mu = 1$ by induction on α since $S_0^\mu = 1$ for any $\mu \in \mathbb{N}$ ■

6.2. Quantum Weyl group action on singular vectors. Let E, F, H be the standard generators of $U_{\hbar} \mathfrak{sl}_2$.

Lemma 6.3. *The following holds in $\mathcal{S}_{\hbar}(\mathcal{M}_{k,2}^*)$,*

$$E v_i^{\mu_1, \mu_2} = [\mu_2 - i] v_i^{\mu_1 + 1, \mu_2 - 1} \quad (6.19)$$

$$F v_i^{\mu_1, \mu_2} = [\mu_1 - i] v_i^{\mu_1 - 1, \mu_2 + 1} \quad (6.20)$$

$$H v_i^{\mu_1, \mu_2} = (\mu_1 - \mu_2) v_i^{\mu_1, \mu_2} \quad (6.21)$$

PROOF. By (5.21),

$$\begin{aligned} E v_i^{\mu_1, \mu_2} &= \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} [\mu_2 - a] e^{\hbar(i - 2a)} X_{11}^{\mu_1 - i + a + 1} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a - 1} X_{22}^a \\ &\quad + \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} [a] X_{11}^{\mu_1 - i + a} X_{21}^{i - a + 1} \otimes X_{12}^{\mu_2 - a} X_{22}^{a - 1} \\ &= \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} [\mu_2 - a] e^{\hbar(i - 2a)} X_{11}^{\mu_1 - i + a + 1} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a - 1} X_{22}^a \\ &\quad - \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar(a + 1)(\mu_2 - a)} [i - a] X_{11}^{\mu_1 - i + a + 1} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a - 1} X_{22}^a \\ &= [\mu_2 - i] \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a)} X_{11}^{\mu_1 + 1 - i + a} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - 1 - a} X_{22}^a \\ &= [\mu_2 - i] v_i^{\mu_1 + 1, \mu_2 - 1} \end{aligned} \quad (6.22)$$

Similarly, by (5.22),

$$\begin{aligned}
F v_i^{\mu_1, \mu_2} &= \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} [\mu_1 - i + a] X_{11}^{\mu_1 - i + a - 1} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a + 1} X_{22}^a \\
&+ \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} [i - a] e^{-\hbar(\mu_1 - \mu_2 - i + 2a)} X_{11}^{\mu_1 - i + a} X_{21}^{i - a - 1} \otimes X_{12}^{\mu_2 - a} X_{22}^{a + 1} \\
&= \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 1)} [\mu_1 - i + a] X_{11}^{\mu_1 - i + a - 1} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a + 1} X_{22}^a \\
&- \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar(a-1)(\mu_2 - a + 2)} [a] e^{-\hbar(\mu_1 - \mu_2 - i + 2a - 2)} X_{11}^{\mu_1 - i + a - 1} X_{21}^{i - a} \otimes X_{12}^{\mu_2 - a + 1} X_{22}^a \\
&= [\mu_1 - i] \sum_{a=0}^i (-1)^a \begin{bmatrix} i \\ a \end{bmatrix} e^{\hbar a(\mu_2 - a + 2)} X_{11}^{\mu_1 - 1 - i + a} X_{21}^{i - a} \otimes X_{12}^{\mu_2 + 1 - a} X_{22}^a \\
&= [\mu_1 - i] v_i^{\mu_1 - 1, \mu_2 + 1}
\end{aligned} \tag{6.23}$$

Finally, (6.21) follows from (5.20) ■

Let now

$$S = \exp_{q^{-1}}(q^{-1} E q^{-H}) \exp_{q^{-1}}(-F) \exp_{q^{-1}}(q E q^H) q^{H(H+1)/2} \tag{6.24}$$

be the generator of the quantum Weyl group of $U_{\hbar} \mathfrak{sl}_2$ ([Lu3, KR, So], we use the form given in [Ka, Sa]) where $q = e^{\hbar}$ and the q -exponential is defined by (6.12).

Proposition 6.4. *The following holds in $\mathcal{S}_{\hbar}(\mathcal{M}_{k,2}^*)$,*

$$S v_i^{\mu_1, \mu_2} = (-1)^{\mu_1 - i} q^{(\mu_1 - i)(\mu_2 - i + 1)} v_i^{\mu_2, \mu_1} \tag{6.25}$$

PROOF. Fix $\mu, i \in \mathbb{N}$ with $2i \leq \mu$. By lemma 6.3, the vectors $u_k = v_i^{\mu - i - k, i + k}$, with $0 \leq k \leq \mu - 2i$ satisfy

$$E u_k = [k] u_{k-1} \tag{6.26}$$

$$F u_k = [\mu - 2i - k] u_{k+1} \tag{6.27}$$

$$H u_k = (\mu - 2i - 2k) u_k \tag{6.28}$$

and therefore span the indecomposable $U_{\hbar} \mathfrak{sl}_2$ -module of dimension $\mu - 2i + 1$. Moreover,

$$u_k = \frac{[\mu - 2i - k]!}{[\mu - 2i]!} F^k u_0 = \frac{[\mu - 2i - k]!}{[\mu - 2i]!} E^{\mu - 2i - k} u_{\mu - 2i} \tag{6.29}$$

Since $\text{Ad}(S)H = -H$, $S u_0$ is proportional to $u_{\mu - 2i}$ and, by (6.24) is therefore equal to

$$(-1)^{\mu - 2i} q^{\mu - 2i} \frac{F^{\mu - 2i}}{[\mu - 2i]!} u_0 = (-1)^{\mu} q^{\mu - 2i} u_{\mu - 2i} \tag{6.30}$$

Next, using $\text{Ad}(S)F = -q^{-H}E$ and (6.29), we find

$$S u_k = \frac{[\mu - 2i - k]!}{[\mu - 2i]!} \text{Ad}(S)F^k S u_0 = (-1)^{\mu - k} q^{(k+1)(\mu - 2i - k)} u_{\mu - 2i - k} \tag{6.31}$$

Thus, setting $\mu = \mu_1 + \mu_2$, so that $v_i^{\mu_1, \mu_2} = u_{\mu_2 - i}$, we find

$$S v_i^{\mu_1, \mu_2} = (-1)^{\mu_1 - i} q^{(\mu_1 - i)(\mu_2 - i + 1)} v_i^{\mu_2, \mu_1} \tag{6.32}$$

as claimed ■

6.3. Identification of $\mathbf{R}$ and quantum Weyl group actions. Fix $1 \leq j \leq n$ and let $R_j^\vee$ be the universal R -matrix of $U_{\hbar} \mathfrak{gl}_k$ acting on the j and $j+1$ tensor copies of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n}$. Let S_j be the element of the quantum Weyl group of $U_{\hbar} \mathfrak{gl}_n$ corresponding to the simple root $\theta_j - \theta_{j+1}$.

Theorem 6.5. *The following holds on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$*

$$R_j^\vee = S_j \cdot e^{-\hbar(D_j^{(n)} + D_j^{(n)} D_{j+1}^{(n)}/k)} \cdot e^{i\pi D_j^{(n)}} \quad (6.33)$$

PROOF. Let $U_{\hbar} \mathfrak{gl}_2^j \subset U_{\hbar} \mathfrak{gl}_n$ be the Hopf subalgebra generated by $E_j^{(n)}, F_j^{(n)}, D_j^{(n)}, D_{j+1}^{(n)}$. By (5.20)–(5.22), $U_{\hbar} \mathfrak{gl}_2^j$ only acts upon the variables X_{ij}, X_{ij+1} , $1 \leq i \leq k$. Thus, $U_{\hbar} \mathfrak{gl}_2^j$ and therefore S_j only act on the j and $j+1$ tensor copies $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)_j, \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)_{j+1}$ of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)^{\otimes n} \cong \mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$. Since this is also the case of $R_j^\vee$, the proof of (6.33) reduces to a computation on the $U_{\hbar} \mathfrak{gl}_k \otimes U_{\hbar} \mathfrak{gl}_2^j$ -module $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)_j \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)_{j+1} \cong \mathcal{S}_{\hbar}(\mathcal{M}_{k,2}^*)$. Both sides of (6.33) clearly commute with $U_{\hbar} \mathfrak{gl}_k$, so it suffices to check their equality on the singular vectors of

$$\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)_j \otimes \mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)_{j+1} = \bigoplus_{\mu_1, \mu_2 \in \mathbb{N}} S_{\hbar}^{\mu_1} \mathbb{C}^k \otimes S_{\hbar}^{\mu_2} \mathbb{C}^k \quad (6.34)$$

i.e., on the vectors $v_i^{\mu_1, \mu_2} \in S_{\hbar}^{\mu_1} \mathbb{C}^k \otimes S_{\hbar}^{\mu_2} \mathbb{C}^k$ of lemma 6.1. By propositions 6.2 and 6.4,

$$R_j^\vee v_i^{\mu_1, \mu_2} = S_j \cdot e^{-\hbar(\mu_1 + \mu_1 \mu_2 / k)} (-1)^{\mu_1} v_i^{\mu_1, \mu_2} \quad (6.35)$$

whence (6.33) since, for any $1 \leq l \leq n$, $D_l^{(n)}$ gives the $\mathbb{N}$ -grading on $\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*)_l$ ■

REMARK. The coincidence of the two representations of B_n studied above was also noted by Baumann [Ba, prop. 12] in the special case when both actions are restricted to the subspace $S_{\hbar}^{\mu_1} \mathbb{C}^k \otimes \cdots \otimes S_{\hbar}^{\mu_n} \mathbb{C}^k$ of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*)$ where $\mu_1 = \cdots = \mu_n = 1$.

7. MONODROMIC REALISATION OF QUANTUM WEYL GROUP OPERATORS

The following is the main result of this paper. It was conjectured for any simple Lie algebra $\mathfrak{g}$ by De Concini around 1995 (unpublished) and, independently, in [TL]. We prove it here for $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$.

Theorem 7.1. *Let $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$, V a finite-dimensional $\mathfrak{g}$ -module, μ a weight of V and*

$$V^\mu = \bigoplus_{\nu \in W\mu} V[\nu] \quad (7.1)$$

the direct sum of the weight spaces of V corresponding to the Weyl group orbit of μ . Let $\pi_\kappa^h : B_{\mathfrak{g}} \rightarrow GL(V^\mu[[h]])$ be the monodromy representation of the connection

$$d - h \sum_{\alpha > 0} \frac{d\alpha}{\alpha} \pi_V(\kappa_\alpha) \quad (7.2)$$

obtained by regarding h as a formal variable. Let $\pi_W : B_{\mathfrak{g}} \rightarrow GL(V^\mu[[h]])$ be the quantum Weyl group action obtained by regarding $V[[h]]$ as a $U_{\hbar} \mathfrak{g}$ -module with $\hbar = 2\pi i h$. Then, π_κ^h and π_W are equivalent.

PROOF. We may assume that V is irreducible with highest weight $\lambda = (\lambda_1, \dots, \lambda_n)$ where $\lambda_i \in \mathbb{N}$. Regard V as a $\mathfrak{gl}_n$ -module by letting $1^{(n)} = \sum_{j=1}^n E_{jj}^{(n)}$ act as multiplication by $|\lambda| = \sum_{i=1}^n \lambda_i$ and fix some $k \geq n$. By lemma 3.4, $V[\nu]$ is isomorphic to the space M_λ^ν of singular vectors of weight λ for the diagonal action of $\mathfrak{gl}_k$ on

$$\mathcal{S}^\nu \mathbb{C}^k = \mathcal{S}^{\nu_1} \mathbb{C}^k \otimes \dots \otimes \mathcal{S}^{\nu_n} \mathbb{C}^k \subset \mathcal{S}(\mathcal{M}_{k,n}^*) \quad (7.3)$$

and, by corollary 3.6, the monodromy representation of the Knizhnik–Zamolodchikov connection (3.17) on $\bigoplus_{\nu \in \mathfrak{S}_k} M_\lambda^\nu$ and π_κ^h are related by

$$\pi_{\text{KZ}}^{\bar{h}}(T_j) = \pi_\kappa^h(T_j) e^{-\pi i h (E_{jj}^{(n)} + E_{j+1j+1}^{(n)} + 2E_{jj}^{(n)} E_{j+1j+1}^{(n)}/k)} e^{i\pi E_{jj}^{(n)}} \quad (7.4)$$

where $\bar{h} = 2h$.

We shall now use the Kohno–Drinfeld theorem to relate $\pi_{\text{KZ}}^{\bar{h}}$ to the R -matrix representation of B_n corresponding to the action of $U_{\hbar} \mathfrak{gl}_k$ on $\mathcal{S}^\nu \mathbb{C}^k[[\hbar]]$. Let for this purpose $\phi : U_{\hbar} \mathfrak{gl}_k \rightarrow U \mathfrak{gl}_k[[\hbar]]$ be an algebra isomorphism whose reduction mod $\hbar$ is the identity and which acts as the identity on the Cartan subalgebras *i.e.*, $\phi(D_i^{(k)}) = E_{ii}^{(k)}$, $1 \leq i \leq k$ [Dr2, prop. 4.3]. Then, $U_{\hbar} \mathfrak{gl}_k$ act on each $\mathcal{S}^{\nu_j} \mathbb{C}^k[[\hbar]]$ via ϕ and on

$$\mathcal{S}^{\nu_1} \mathbb{C}^k[[\hbar]] \otimes \dots \otimes \mathcal{S}^{\nu_n} \mathbb{C}^k[[\hbar]] = \mathcal{S}^{\nu_1} \mathbb{C}^k \otimes \dots \otimes \mathcal{S}^{\nu_n} \mathbb{C}^k[[\hbar]] = \mathcal{S}^\nu \mathbb{C}^k[[\hbar]] \quad (7.5)$$

via the n -fold coproduct $\Delta^{(n)} : U_{\hbar} \mathfrak{gl}_k \rightarrow U_{\hbar} \mathfrak{gl}_k^{\otimes n}$ recursively defined by

$$\Delta^{(1)} = \text{id} \quad (7.6)$$

$$\Delta^{(a+1)} = \Delta \otimes \text{id}^{\otimes(a-1)} \circ \Delta^{(a)}, \quad a \geq 1 \quad (7.7)$$

where $\Delta = \Delta^{(2)}$ is given by (4.9)–(4.11).

Let Δ_0 be the standard, cocommutative coproduct on $U \mathfrak{gl}_k$ so that $U \mathfrak{gl}_k$ acts on $\mathcal{S}^\nu \mathbb{C}^k$ via $\Delta_0^{(n)} : U \mathfrak{gl}_k \rightarrow U \mathfrak{gl}_k^{\otimes n}$ defined as in (7.6)–(7.7) with Δ replaced by Δ_0 . Since $\Delta = \Delta_0 + o(\hbar)$ and $H^1(\mathfrak{sl}_k, \mathfrak{sl}_k \oplus \mathfrak{sl}_k) = 0$, there exists a twist $F = 1 \otimes 1 + o(\hbar) \in U \mathfrak{gl}_k^{\otimes(2)}[[\hbar]]$ such that, for any $x \in U_{\hbar} \mathfrak{gl}_k$,

$$\phi \otimes \phi \circ \Delta(x) = F \Delta_0(\phi(x)) F^{-1} \quad (7.8)$$

It follows that the actions of $U_{\hbar} \mathfrak{gl}_k$ and $U \mathfrak{gl}_k[[\hbar]]$ on $\mathcal{S}^\nu \mathbb{C}^k$ are related by

$$\phi^{\otimes(n)} \circ \Delta^{(n)}(x) = F^{(n)} \Delta_0^{(n)}(\phi(x)) F^{(n)-1}, \quad x \in U_{\hbar} \mathfrak{gl}_k \quad (7.9)$$

where $F^{(n)} \in U \mathfrak{gl}_k^{\otimes n}[[\hbar]]$ is recursively defined by

$$F^{(1)} = 1 \quad (7.10)$$

$$F^{(a+1)} = F \otimes 1^{\otimes(a-1)} \cdot \Delta_0 \otimes \text{id}^{\otimes(a-1)}(F^{(a)}), \quad a \geq 1 \quad (7.11)$$

Let now $M_\lambda^\nu \subset \mathcal{S}^\nu \mathbb{C}^k$ and $M_{\hbar,\lambda}^\nu \subset \mathcal{S}^\nu \mathbb{C}^k[[\hbar]]$ be the subspaces of vectors of highest weight λ for the actions of $\mathfrak{gl}_k$ and $U_{\hbar} \mathfrak{gl}_k$ respectively. We shall need the following

Lemma 7.2. $F^{(n)} M_\lambda^\nu[[\hbar]] = M_{\hbar,\lambda}^\nu$.

PROOF. Let $\mathcal{S}^\nu \mathbb{C}^k(\lambda) \subset \mathcal{S}^\nu \mathbb{C}^k$ and $\mathcal{S}^\nu \mathbb{C}^k[[\hbar]](\lambda) \subset \mathcal{S}^\nu \mathbb{C}^k[[\hbar]]$ be the isotypical components of types V_λ and $V_\lambda[[\hbar]]$ for $\mathfrak{gl}_k$ and $U_{\hbar} \mathfrak{gl}_k$ respectively. The subspace $F^{(n)} \mathcal{S}^\nu \mathbb{C}^k(\lambda)[[\hbar]]$ is invariant under $U_{\hbar} \mathfrak{gl}_k$ by (7.9) and its reduction mod $\hbar$ is equal to $\mathcal{S}^\nu \mathbb{C}^k(\lambda)$ since $F^{(n)} = 1^{\otimes(n)} + o(\hbar)$. Thus, by proposition 4.1, $F^{(n)} \mathcal{S}^\nu \mathbb{C}^k(\lambda)[[\hbar]]$ is isomorphic to $\mathcal{S}^\nu \mathbb{C}^k(\lambda)[[\hbar]]$ and therefore contained in $\mathcal{S}^\nu \mathbb{C}^k[[\hbar]](\lambda)$. Since this holds for any λ , the inclusion is an equality. Noting

now that $F^{(n)}$ is equivariant for the action of the Cartan subalgebras of $\mathfrak{gl}_k$ and $U_{\hbar}\mathfrak{gl}_k$ since $\Delta_0 \circ \phi(h) = \phi \otimes \phi \circ \Delta(h)$ for any $h \in \mathfrak{h}$, we get that

$$\begin{aligned} F^{(n)} M_{\lambda}^{\nu}[\hbar] &= F^{(n)} \left(\mathcal{S}^{\nu} \mathbb{C}^k(\lambda)[\lambda][\hbar] \right) \\ &= F^{(n)} \left(\mathcal{S}^{\nu} \mathbb{C}^k(\lambda)[\hbar] \right) [\lambda] \\ &= \mathcal{S}^{\nu} \mathbb{C}^k[\hbar](\lambda)[\lambda] \\ &= M_{\hbar, \lambda}^{\nu} \end{aligned} \tag{7.12}$$

as claimed ■

Summarising, we have an action of B_n on $\bigoplus_{\nu \in \mathfrak{S}_n \mu} M_{\lambda}^{\nu}$ via the monodromy of the Knizhnik–Zamolodchikov connection and an action of B_n on $\bigoplus_{\nu \in \mathfrak{S}_n \mu} M_{\hbar, \lambda}^{\nu}$ via the R -matrix representation of $U_{\hbar}\mathfrak{sl}_k$. Drinfeld’s theorem [Dr3, Dr4, Dr5] asserts that the twist F may be chosen so that

$$F^{(n)} : \bigoplus_{\nu \in \mathfrak{S}_n \mu} M_{\lambda}^{\nu}[\hbar] \longrightarrow \bigoplus_{\nu \in \mathfrak{S}_n \mu} M_{\hbar, \lambda}^{\nu} \tag{7.13}$$

is B_n -equivariant so that, for any $1 \leq j \leq n-1$,

$$\pi_{\text{KZ}}^{\hbar}(T_j) = F^{(n)-1} R_j^{\vee} F^{(n)} \tag{7.14}$$

where $R_j^{\vee}$ is the universal R -matrix for $U_{\hbar}\mathfrak{sl}_k$ acting on the j and $j+1$ copies of $\bigoplus_{\nu \in \mathfrak{S}_n \mu} \mathcal{S}^{\nu} \mathbb{C}^k[\hbar]$ and $\hbar = \pi i \bar{\hbar}$.

Let now $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \cong (\mathcal{S}_{\hbar}(\mathcal{M}_{k,1}^*))^{\otimes n}$ be the algebra of functions on quantum $k \times n$ matrix space defined in section 5 and, for any $\nu \in \mathbb{N}^n$, let

$$\mathcal{S}_{\hbar}^{\nu} \mathbb{C}^k = \mathcal{S}_{\hbar}^{\nu_1} \mathbb{C}^k \otimes \cdots \otimes \mathcal{S}_{\hbar}^{\nu_n} \mathbb{C}^k \tag{7.15}$$

be the space of polynomials which are homogeneous of degree ν_j in the variables $X_{1j}, \dots, X_{kj}$, for any $1 \leq j \leq n$. By (5.17)–(5.19), $\mathcal{S}_{\hbar}^{\nu} \mathbb{C}^k / \hbar \mathcal{S}_{\hbar}^{\nu} \mathbb{C}^k$ is the $\mathfrak{gl}_k$ -module $\mathcal{S}^{\nu} \mathbb{C}^k$ so that, by proposition 4.1, we may identify $\mathcal{S}_{\hbar}^{\nu} \mathbb{C}^k$ with $\mathcal{S}^{\nu} \mathbb{C}^k[\hbar]$ as $U_{\hbar}\mathfrak{gl}_k$ -modules.

By theorem 5.5 and the fact that the $\mathbb{N}$ -grading on the j th tensor factor of $\mathcal{S}_{\hbar}(\mathcal{M}_{k,n}^*) \cong (\mathcal{S}_{\hbar}(\mathcal{M}_{1,k}^*))^{\otimes n}$ is given by the action of $D_j^{(n)}$, the space $M_{\hbar, \lambda}^{\nu}$ is isomorphic to the subspace $V_{\lambda}^{(n)}[\nu][\hbar] \subset V_{\lambda}^{(n)}[\hbar]$ of weight ν . Using now theorem 6.5, we find

$$\begin{aligned} \pi_{\kappa}^{\hbar}(T_j) &= F^{(n)-1} \cdot S_j \cdot e^{-\hbar(D_j^{(n)} + D_j^{(n)} D_{j+1}^{(n)} / k)} \cdot e^{i\pi D_j^{(n)}} \cdot F^{(n)} \\ &\quad \cdot e^{\pi i \hbar (E_j^{(n)} + E_{j+1}^{(n)} + 2E_j^{(n)} E_{j+1}^{(n)} / k)} \cdot e^{i\pi E_j^{(n)}} \end{aligned} \tag{7.16}$$

where $S_j = \pi_W(T_j)$ is the quantum Weyl group element of $U_{\hbar}\mathfrak{gl}_n$ corresponding to the simple root $\alpha_j = \theta_j - \theta_{j+1}$. Since for any l , $E_l^{(n)} = D_l^{(n)}$ on $V^{\mu} \cong \bigoplus_{\nu \in \mathfrak{S}_k} M_{\lambda}^{\nu}$ we get

$$\begin{aligned} \pi_{\kappa}^{\hbar}(T_j) &= F^{(n)-1} S_j e^{-\hbar/2(D_j^{(n)} - D_{j+1}^{(n)})} F^{(n)} \\ &= F^{(n)-1} S_j e^{-\hbar H_j^{(n)}/2} F^{(n)} \\ &= F^{(n)-1} e^{\hbar \rho^{(n)}/2} S_j e^{-\hbar \rho^{(n)}/2} F^{(n)} \end{aligned} \tag{7.17}$$

where $\rho^{(n)} = \frac{1}{2} \sum_{j=1}^n (n-2j+1) D_j^{(n)}$ is the half-sum of the positive (co)roots of $\mathfrak{sl}_n$ ■

REMARK. Theorem 7.1 is proved in [TL] for the following pairs $(\mathfrak{g}, V)$ where $\mathfrak{g}$ is a simple Lie algebra and V an irreducible, finite-dimensional representation

1. $\mathfrak{g} = \mathfrak{sl}_2$ and V is any irreducible representation.
2. $\mathfrak{g} = \mathfrak{sl}_n$ and V is a fundamental representation.
3. $\mathfrak{g} = \mathfrak{so}_n$ and V is the vector or a spin representation.
4. $\mathfrak{g} = \mathfrak{sp}_n$ and V is the defining vector representation.
5. $\mathfrak{g} = \mathfrak{e}_6, \mathfrak{e}_7$ and V is a minuscule representation.
6. $\mathfrak{g} = \mathfrak{g}_2$ and V is the 7-dimensional representation.
7. $\mathfrak{g}$ is any simple Lie algebra and $V \cong \mathfrak{g}$ its adjoint representation.

REFERENCES

- [Ba] P. Baumann, *The q -Weyl Group of a q -Schur Algebra*, preprint, 1999.
- [Br] E. Brieskorn, *Die Fundamentalgruppe des Raumes der Regulären Orbits einer Endlichen Komplexen Spiegelungsgruppe*, Invent. Math. **12** (1971), 57–61.
- [CP] V. Chari, A. Pressley, *A Guide to Quantum Groups*. Cambridge University Press, Cambridge, 1994.
- [Dr1] V. G. Drinfeld, *Quantum Groups*, Proceedings of the International Congress of Mathematicians, Berkeley 1986, 798–820.
- [Dr2] V. G. Drinfeld, *On Almost Cocommutative Hopf Algebras*, Leningrad Math. J. **1** (1990), 321–342.
- [Dr3] V. G. Drinfeld, *Quasi-Hopf Algebras*, Leningrad Math. J. **1** (1990), 1419–57.
- [Dr4] V. G. Drinfeld, *On Quasitriangular Quasi-Hopf Algebras and on a Group that is closely connected with $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$* , Leningrad Math. J. **2** (1991), 829–860.
- [Dr5] V. G. Drinfeld, *On the Structure of Quasitriangular Quasi-Hopf Algebras*, Functional Anal. Appl. **26** (1992), 63–65.
- [FMTV] G. Felder, Y. Markov, V. Tarasov, A. Varchenko, *Differential Equations Compatible with KZ Equations*, Math. Phys. Anal. Geom. **3** (2000), 139–177.
- [Ga] F. Galdi, Ph.D. dissertation, Università di Roma II "Tor Vergata", 1995.
- [GS] V. Guillemin, S. Sternberg, *The Gel'fand-Cetlin System and Quantization of the Complex Flag Manifolds*, J. Funct. Anal. **52** (1983), 106–128.
- [HK] J.-C. Hausmann, A. Knutson, *Polygon Spaces and Grassmannians*. Enseign. Math. **43** (1997), 173–198.
- [Ho] R. Howe, *Remarks on Classical Invariant Theory*, Trans. Amer. Math. Soc. **313** (1989), 539–570.
- [Ji1] M. Jimbo, *A q -Difference Analogue of $U(\mathfrak{g})$ and the Yang-Baxter Equation*. Lett. Math. Phys. **10** (1985), 63–69.
- [Ji2] M. Jimbo, *A q -Analogue of $U\mathfrak{gl}(N+1)$, Hecke Algebra, and the Yang-Baxter Equation*, Lett. Math. Phys. **11** (1986), 247–252.
- [KM] M. Kapovich, J. Millson, *The Symplectic Geometry of Polygons in Euclidean Space*, J. Differential Geom. **44** (1996), 479–513.
- [Ka] M. Kashiwara, *Private Note on Finite-Dimensional Representations of Quantized Affine Algebras*, unpublished notes.
- [KR] A. N. Kirillov, N. Reshetikhin, *q -Weyl Group and a Multiplicative Formula for Universal R -Matrices*, Comm. Math. Phys. **134** (1990), 421–431.
- [Kn] A. Knutson, personal communication to J. Millson, May 1999.
- [Ko1] T. Kohno, *Quantized Enveloping Algebras and Monodromy of Braid Groups*, preprint, 1988.
- [Ko2] T. Kohno, *Integrable Connections Related to Manin and Schechtman's Higher Braid Groups*, Illinois J. Math. **34** (1990), 476–484.
- [LS] S. Z. Levendorskii, Y. S. Soibelman, *Some applications of the Quantum Weyl Groups*, J. Geom. Phys. **7** (1990), 241–254.
- [Lu1] G. Lusztig, *On Quantum Groups*, J. Algebra **131** (1990), 466–475.
- [Lu2] G. Lusztig, *Finite-Dimensional Hopf Algebras Arising from Quantized Universal Enveloping Algebra*. J. Amer. Math. Soc. **3** (1990), 257–296.
- [Lu3] G. Lusztig, *Introduction to Quantum Groups*. Progress in Mathematics, 110. Birkhäuser Boston, 1993.
- [Mc] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*. 2nd edition, Oxford University Press, 1995.
- [MTL] J. Millson, V. Toledano Laredo, *Casimir Operators and Monodromy Representations of Generalised Braid Groups*, in preparation.
- [PW] B. Parshall, J.-P. Wang, *Quantum Linear Groups*. Mem. Amer. Math. Soc. **89** (1991), n. 439.

- [Ro] M. Rosso, *An analogue of P.B.W. Theorem and the Universal R-Matrix for $U_{\hbar}\mathfrak{sl}(N+1)$* . Comm. Math. Phys. **124** (1989), 307–318.
- [Sa] Y. Saito, *PBW Basis of Quantized Universal Enveloping Algebras*, Publ. Res. Inst. Math. Sci. **30** (1994), 209–232.
- [So] Y. S. Soibelman, *Algebra of Functions on a Compact Quantum Group and its Representations*, Leningrad Math. J. **2** (1991), 161–178.
- [TV] V. Tarasov, A. Varchenko, *Difference Equations Compatible with Trigonometric KZ Differential Equations*, Internat. Math. Res. Notices **2000**, 801–829.
- [Ti] J. Tits, *Normalisateurs de Tores. I. Groupes de Coxeter Etendus*, J. Algebra **4** (1966), 96–116.
- [TL] V. Toledano Laredo, *Monodromy Representations of Generalised Braid Groups and Quantum Weyl Groups*, in preparation.
- [Zh] D. P. Zhelobenko, *Compact Lie Groups and their Representations.*, Translations of Mathematical Monographs, Vol. 40. American Mathematical Society, 1973.

MSRI
 1000 CENTENNIAL DRIVE
 BERKELEY, CA 94720–5070
 TOLEDANO@MSRI.ORG

PERMANENT ADDRESS :

INSTITUT DE MATHÉMATIQUES DE JUSSIEU
 UMR 7586, CASE 191
 UNIVERSITÉ PIERRE ET MARIE CURIE
 4, PLACE JUSSIEU
 F–75252 PARIS CEDEX 05
 TOLEDANO@MATH.JUSSIEU.FR